

Erratum

doi: 10.5123/S1679-49742018000200026

In the article “Attendance of children and adolescents with psychoactive substance use disorders performed by Psychosocial Care Centers in Brazil, 2008-2012”, DOI: 10.5123/S1679-49742018000200002, published on *Epidemiology and Health Services*, 27(2):1-12, pages 4,5 and 6:

Original text:

The ethnicity/skin color white was indicated in 29.1% of records, followed by brown, with 19.6%, and black, with 4.8%.

Table 1 – Demographic Characteristics and nosological of attendance for children and adolescents with mental and behavioral disorders by use of psychoactive substances performed in Psychosocial Care Centers, Brazil, 2008-2012

Variables	Year										Total	
	2008		2009		2010		2011		2012			
	n	%	n	%	n	%	n	%	n	%	n	%
Sex												
Female	3,941	19.4	4,131	17.5	5,490	17.8	6,755	18.1	8,081	20.7	28,398	18.8
Male	16,501	80.7	19,531	82.5	25,412	82.2	30,580	81.9	30,908	79.3	122,932	81.2
Age group (in years)												
5-9	256	1.2	201	0.8	219	0.7	298	0.8	213	0.5	1,187	0.8
10-14	3,540	17.3	3,879	16.5	4,783	15.5	5,200	13.9	5,608	14.5	23,010	15.3
15-19	16,646	81.4	19,582	2.8	25,900	83.8	31,837	85.3	33,168	5.1	127,133	4.0
Ethnicity/skin color												
White	6,835	33.4	7,105	30.0	8,083	26.2	10,434	27.9	11,584	29.7	44,041	29.1
Black	1,136	5.6	1,145	4.8	1,482	4.8	1,741	4.7	1,828	4.7	7,332	4.8
Brown	4,245	20.8	4,353	18.4	5,373	17.4	7,381	19.8	8,268	21.2	29,620	19.6
Asian	274	1.3	284	1.2	264	0.8	261	0.7	448	1.1	1,531	1.0
Indigenous	9	0.0	5	0.0	3	0.0	22	0.1	30	0.1	69	0.0
No information	7,943	38.9	10,770	45.5	15,697	50.8	17,496	46.9	16,831	43.2	68,737	45.4
Cause of attendance												
F10 ^a (alcohol)	2,242	11.0	2,620	11.1	2,699	8.7	2,849	7.6	3,207	8.2	13,617	9.0
F11 ^b (opioid)	80	0.4	76	0.3	86	0.3	123	0.3	182	0.5	547	0.4
F12 ^c (cannabinoids)	2,924	14.3	3,600	15.2	4,455	14.4	5,870	15.7	6,729	17.2	23,578	15.6
F13 ^d (sedatives)	62	0.3	51	0.2	92	0.3	120	0.3	53	0.1	378	0.2
F14 ^e (cocaine)	2,511	12.3	3,284	13.9	5,145	16.6	6,721	18.0	6,026	15.5	23,687	15.6
F15 ^f (stimulants)	150	0.7	49	0.2	71	0.2	151	0.4	181	0.5	602	0.4
F16 ^g (hallucinogens)	29	0.1	50	0.2	51	0.2	226	0.6	70	0.2	426	0.3
F17 ^h (tobacco)	258	1.3	186	0.8	262	0.8	312	0.8	321	0.8	1,339	0.9
F18 ⁱ (solvents)	470	2.3	294	1.2	255	0.8	197	0.5	161	0.4	1,377	0.9
F19 ^j (multiples SPA ^k)	11,716	57.3	13,452	56.8	17,786	57.6	20,766	55.6	22,059	56.6	85,779	56.7
Total	20,442	13.5	23,662	15.6	30,902	20.4	37,335	24.7	39,119	25.8	151,330	100.0

a) F10: Mental and behavioral disorders due to use of alcohol.

b) F11: Mental and behavioural disorders due to use of opioids.

c) F12: Mental and behavioral disorders due to use of cannabinoids.

d) F13: Mental and behavioral disorders due to the use of sedatives and hypnotics.

e) F14: Mental and behavioral disorders due to use of cocaine.

f) F15: Mental and behavioral disorders due to use of other stimulants, including caffeine.

g) F16: Mental and behavioral disorders due to the use of hallucinogens.

h) F17: Mental and behavioral disorders due to use of tobacco.

i) F18: Mental and behavioral disorders due to use of volatile solvents.

j) F19: Mental and behavioral disorders due to use of multiple drugs and the use of other psychoactive substances.

k) SPA: Psychoactive Substances.

Sources: Authorization for outpatient procedures (APAC); International Statistical Classification of Diseases and Related Health Problems, 10th Revision (ICD-10).

Table 2 – Distribution of records of attendance due to mental and behavioral disorders by use of psychoactive substances, according to type of Psychosocial Care Center and great national region, Brazil, 2008-2012

Type of CAPS	Region										Total	
	North		Northeast		Southeast		South		Midwest			
	n	%	n	%	n (63,165)	%	n	%	n	%	n	%
CAPS I ^a	210	5.7	2,098	7.4	981	1.5	2,416	5.3	307	2.8	6,012	4.0
CAPS II ^b	98	2.7	417	1.5	4,645	7.3	215	0.5	7	0.1	5,382	3.6
CAPS III ^c	6	0.2	176	0.6	222	0.3	17	0.0	95	0.9	516	0.3
CAPSi ^d	39	1.7	2,664	9.4	6,380	10.1	6,482	14.3	129	1.2	15,694	10.4
CAPS AD ^e / CAPS AD III ^f	3,322	90.4	22,973	81.1	50,937	80.6	36,245	79.9	10,249	95.0	123,726	81.8
Total	3,975	100.0	28,328	100.0	63,165	100.0	45,375	100.0	10,787	100.0	151,330	100.0

a) CAPS I: Psychosocial Care Center type I.

b) CAPS II: Psychosocial Care Center type II.

c) CAPS III: Psychosocial Care Center type III.

d) CAPSi: Psychosocial Care Center for Childhood and Youth.

e) CAPS AD: Psychosocial Care Center for Alcohol and other Drugs.

f) CAPS AD III: Psychosocial Care Center for Alcohol and other Drugs 24 Hours.

Sources: Authorization for outpatient procedures (APAC); National Database on Health Units (CNES).

Corrected text:

The ethnicity/skin color white was indicated in 29.1% of records, followed by brown, with 19.6%, and black, with 4.9%.

Table 1 – Demographic Characteristics and nosological of attendance for children and adolescents with mental and behavioral disorders by use of psychoactive substances performed in Psychosocial Care Centers, Brazil, 2008-2012

Variables	Year										Total	
	2008		2009		2010		2011		2012			
	n	%	n	%	n	%	n	%	n	%	n	%
Sex												
Female	3,941	19.3	4,131	17.5	5,490	17.8	6,755	18.1	8,081	20.7	28,398	18.8
Male	16,501	80.7	19,531	82.5	25,412	82.2	30,580	81.9	30,908	79.3	122,932	81.2
Age group (in years)												
5-9	256	1.3	201	0.8	219	0.7	298	0.8	213	0.5	1,187	0.8
10-14	3,540	17.3	3,879	16.5	4,783	15.5	5,200	13.9	5,608	14.5	23,010	15.2
15-19	16,646	81.4	19,582	82.7	25,900	83.8	31,837	85.3	33,168	85.0	127,133	84.0
Ethnicity/skin color												
White	6,835	33.4	7,105	30.0	8,083	26.2	10,434	27.9	11,584	29.7	44,041	29.1
Black	1,136	5.6	1,145	4.8	1,482	4.8	1,741	4.7	1,828	4.7	7,332	4.9
Brown	4,245	20.8	4,353	18.4	5,373	17.4	7,381	19.8	8,268	21.2	29,620	19.6
Asian	274	1.3	284	1.2	264	0.8	261	0.7	448	1.1	1,531	1.0
Indigenous	9	0.0	5	0.0	3	0.0	22	0.1	30	0.1	69	0.0
No information	7,943	38.9	10,770	45.6	15,697	50.8	17,496	46.8	16,831	43.2	68,737	45.4
Cause of attendance												
F10 ^a (alcohol)	2,242	11.0	2,620	11.1	2,699	8.7	2,849	7.6	3,207	8.2	13,617	9.0
F11 ^b (opioid)	80	0.4	76	0.3	86	0.3	123	0.3	182	0.5	547	0.4
F12 ^c (cannabinoids)	2,924	14.3	3,600	15.2	4,455	14.4	5,870	15.7	6,729	17.2	23,578	15.6
F13 ^d (sedatives)	62	0.3	51	0.2	92	0.3	120	0.3	53	0.1	378	0.2
F14 ^e (cocaine)	2,511	12.3	3,284	13.9	5,145	16.6	6,721	18.0	6,026	15.5	23,687	15.6
F15 ^f (stimulants)	150	0.7	49	0.2	71	0.2	151	0.4	181	0.5	602	0.4
F16 ^g (hallucinogens)	29	0.1	50	0.2	51	0.2	226	0.6	70	0.2	426	0.3
F17 ^h (tobacco)	258	1.3	186	0.8	262	0.9	312	0.9	321	0.9	1,339	0.9
F18 ⁱ (solvents)	470	2.3	294	1.2	255	0.8	197	0.6	161	0.4	1,377	0.9
F19 ^j (multiples SPA ^k)	11,716	57.3	13,452	56.9	17,786	57.6	20,766	55.6	22,059	56.6	85,779	56.7
Total	20,442	13.5	23,662	15.6	30,902	20.4	37,335	24.7	39,119	25.8	151,330	100.0

a) F10: Mental and behavioral disorders due to use of alcohol.

b) F11: Mental and behavioural disorders due to use of opioids.

c) F12: Mental and behavioral disorders due to use of cannabinoids.

d) F13: Mental and behavioral disorders due to the use of sedatives and hypnotics.

e) F14: Mental and behavioral disorders due to use of cocaine.

f) F15: Mental and behavioral disorders due to use of other stimulants, including caffeine.

g) F16: Mental and behavioral disorders due to the use of hallucinogens.

h) F17: Mental and behavioral disorders due to use of tobacco.

i) F18: Mental and behavioral disorders due to use of volatile solvents.

j) F19: Mental and behavioral disorders due to use of multiple drugs and the use of other psychoactive substances.

k) SPA: Psychoactive Substances.

Sources: Authorization for outpatient procedures (APAC); International Statistical Classification of Diseases and Related Health Problems, 10th Revision (ICD-10).

Table 2 – Distribution of records of attendance due to mental and behavioral disorders by use of psychoactive substances, according to type of Psychosocial Care Center and great national region, Brazil, 2008-2012

Type of CAPS	Region										Total	
	North		Northeast		Southeast		South		Midwest			
	n	%	n	%	n (63,165)	%	n	%	n	%	n	%
CAPS I ^a	210	5.7	2,098	7.4	981	1.6	2,416	5.3	307	2.8	6,012	4.0
CAPS II ^b	98	2.7	417	1.5	4,645	7.4	215	0.5	7	0.1	5,382	3.5
CAPS III ^c	6	0.2	176	0.6	222	0.3	17	0.0	95	0.9	516	0.3
CAPSi ^d	39	1.0	2,664	9.4	6,380	10.1	6,482	14.3	129	1.2	15,694	10.4
CAPS AD ^e / CAPS AD III ^f	3,322	90.4	22,973	81.1	50,937	80.6	36,245	79.9	10,249	95.0	123,726	81.8
Total	3,675	100.0	28,328	100.0	63,165	100.0	45,375	100.0	10,787	100.0	151,330	100.0

a) CAPS I: Psychosocial Care Center type I.

b) CAPS II: Psychosocial Care Center type II.

c) CAPS III: Psychosocial Care Center type III.

d) CAPSi: Psychosocial Care Center for Childhood and Youth.

e) CAPS AD: Psychosocial Care Center for Alcohol and other Drugs.

f) CAPS AD III: Psychosocial Care Center for Alcohol and other Drugs 24 Hours.

Sources: Authorization for outpatient procedures (APAC); National Database on Health Units (CNES).

In the article “Prevalence of health services utilization in Brazil: a systematic review and meta-analysis”, DOI: 10.5123/S1679-49742017000300016, published on *Epidemiology and Health Services*, 26(3):

Original text:

Figure 2 presents the prevalences of medical visits found in the 18 studies,^{17-20,22-25,27,29-34,36,37,44}

Corrected text:

Figure 2 presents the prevalences of medical visits found in the 18 studies,^{17-25, 27, 29-34, 36, 37}

N.B.: The following references should be taken to have one number less than the number shown, i.e. consult the reference immediately prior to the reference cited.

Original text:	Corrected text:
45	44
46	45
47	46
48	47
49	48
50	49
51	50
52	51
53	52
54	53
55	54
56	55
57	56
58	57
59	58
60	59
61	60
62	61
63	62
64	63
65	64
66	65
67	66
68	67
69	68
70	69
71	70
72	71
73	72
74	73
75	74

In the article “Congenital syndrome associated with Zika virus infection among live births in Brazil: a description of the distribution of reported and confirmed cases in 2015-2016”, DOI: 10.5123/S1679-49742018000200014, published on *Epidemiology and Health Services*, 27(2):1-12, page 10:

Original text:

França GVA and Garcia LP participated in the conception and design of the study. França GVA, Garcia LP, I RV, Carmo GMI, loyal MB and Garcia MHO performed the analysis and interpretation of the results, writing and critical review of the manuscript.

Corrected text:

França GVA and Garcia LP participated in the conception and design of the study. França GVA, Garcia LP, Pedi VD, Carmo GMI, Leal MB and Garcia MHO performed the analysis and interpretation of the results, writing and critical review of the manuscript.