

Impact and effectiveness of meningococcal vaccines: a review

Lucia Helena De Oliveira,¹ Barbara Jauregui,¹ Ana Flavia Carvalho,² and Norberto Giglio³

Suggested citation

De Oliveira LH, Jauregui B, Carvalho AF, Giglio N. Impact and effectiveness of meningococcal vaccines: a review. Rev Panam Salud Publica. 2017;41:e158. doi: 10.26633/RPSP.2017.158

ABSTRACT

Objectives. To summarize and critically evaluate the evidence on the impact and effectiveness of meningococcal vaccination programs around the world in order to inform decisionmaking in Latin America and the Caribbean.

Methods. A review of the literature was conducted following several components of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. PubMed Central® was searched for papers published in any language from January 1999 – March 2017.

Results. In all, 32 studies were included, most of which evaluated the meningococcal C conjugate vaccine. Fourteen studies measured effectiveness and 30 measured impact. The effectiveness of polysaccharide vaccines was 65% – 83.7% (different age groups), while the effectiveness of the conjugate vaccines was 66% – 100%. Incidence decline of laboratory-confirmed meningococcal disease for the conjugate vaccine ranged from 77% – 100% among different ages groups. The only study that evaluated the protein subunit vaccine reported a vaccine effectiveness of 82.9%.

Conclusions. The studies reviewed show impact and effectiveness of both polysaccharide vaccines and conjugate vaccines on vaccine-serogroup meningococcal disease. The conjugate vaccines, however, show higher impact and effectiveness with longer-lasting protection over the polysaccharide vaccines. Given the variance in potential use of a meningococcal vaccine, epidemiological surveillance systems should be strengthened to inform national decisions.

Keywords

Meningococcal vaccines; immunization programs; decision making, prevention & control; review; Latin America.

Meningococcal disease is a global problem that occurs in all countries, but its epidemiology varies substantially by capsular group (1). Virulent strains of *Neisseria*

Meningitidis have a polysaccharide capsule, with types A, B, C, W, X, and Y commonly causing invasive infections (2).

The incidence of invasive meningococcal disease (IMD) is highest in children less than 1 year of age and remains relatively high until approximately 5 years. Although the incidence tends to decrease in older children, it usually peaks again during adolescence and young adulthood when individuals are

living in close quarters. Incidence again tapers off in older adults. This pattern has been observed at both regional and country levels and with individual serogroups. The morbidity and mortality from IMD is substantial, with a case-fatality rate close to 10% in developed countries (3). Many survivors experience permanent debilitating sequelae, such as hearing loss, neurologic impairments, or limb loss (4).

¹ Comprehensive Family Immunization Unit, Pan American Health Organization, Regional Office of the World Health Organization, Washington, DC, United States of America. Send correspondence to Lucia Helena De Oliveira, oliveirl@paho.org

² Albert B. Sabin Vaccine Institute, Washington, DC, United States.

³ Hospital Gutierrez, Buenos Aires, Argentina.

 This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs 3.0 IGO License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited. No modifications or commercial use of this article are permitted. In any reproduction of this article there should not be any suggestion that PAHO or this article endorse any specific organization or products. The use of the PAHO logo is not permitted. This notice should be preserved along with the article's original URL.

The development of meningococcal vaccines began in the 1960s with the creation of the polysaccharide vaccines, which are immunogenic in older children and adults, but less so in infants. To cover this younger population, conjugate vaccines were developed in 1999. Polysaccharide vaccines and conjugate vaccines based on the meningococcal capsule are now available against all meningococcal strains related to serogroups A, C, W, and Y. Regarding protection against serogroup B, a broadly protective serogroup B recombinant vaccine has recently been approved in several countries (5, 6).

According to the World Health Organization's (WHO) Vaccine Preventable Diseases Monitoring System, as of 2016, a total of 39 countries had introduced a meningococcal vaccine in their routine immunization schedules. The first country to introduce the conjugate vaccine was the United Kingdom in 1999. Additionally, 30 countries have introduced a meningococcal vaccine for specific high-risk groups only. Infants and young children less than 2 years of age are targeted in 27 countries, while nine countries target children 2 years of age or older and 11 countries target adolescents. Only five countries have introduced a meningococcal vaccine in a geographically restricted area within the country, while the rest have introduced it nationwide (7, 8).

Some countries in Latin America and the Caribbean have decided to implement a nationwide, routine immunization program using meningococcal vaccines. Brazil was the first to introduce a conjugated meningococcal C vaccine (MenC), while Cuba was the first to introduce an outer-membrane protein B vaccine (9). Afterwards, Puerto Rico and Chile introduced the ACWY tetravalent polysaccharide-protein conjugate vaccines in 2008 and 2012, respectively; Argentina did likewise in 2017 (10, 11).

Monitoring post-licensure impact of meningococcal vaccines is important for documenting appropriate gains in terms of expected morbidity and mortality reduction. This information is scarce and scattered in Latin America and the Caribbean, making it difficult for decisionmakers to base policy on evidence. This review aims to summarize the available evidence on the impact and effectiveness of commercially-available vaccines on meningococcal disease, invasive meningococcal disease, and death due to meningococcus

in all age groups, as reported by several countries. The intent is to inform decision making among countries of Latin America and the Caribbean that are considering introducing the meningococcal vaccine.

MATERIALS AND METHODS

Literature search

A literature search following several components of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines⁴ was performed on PubMed Central® (National Library of Medicine, Bethesda, Maryland, United States) to identify all available studies on the effects of meningococcal vaccines on meningococcal disease cases and mortality. The main search terms were: "meningococcal vaccine," "immunization program," "mass vaccination," AND "effectiveness" OR "evaluation" OR "impact" OR "benefits," "hospitalization," "notifications," "decline" OR "reduction." Terms used for exclusion of papers were: "meningococcal vaccines/economics" OR "immunization/economics" or "cost-benefit analysis" OR "MenAfriVac" OR "disease outbreaks" OR "randomized controlled trial." The search included all papers published from 1 January 1999 – 15 March 2017, the day on which the search was conducted.

Inclusion and exclusion criteria

Studies of primary impact and effectiveness of currently available meningococcal vaccines were included, published from January 1999 – March 2017, in any language. The review focused on the meningococcal polysaccharide, polysaccharide-protein conjugate, and protein sub unit vaccines, and considered any serogroup except for serogroup A. The outcomes of interest were clinically-compatible and/or laboratory-confirmed meningococcal cases and deaths due to meningococcal disease.

The following study types were excluded: case series, case reports, and randomized controlled trials; and economic, cost-benefit, and modelling studies.

⁴ An evidence-based minimum set of items for reporting in systematic reviews and meta-analyses, focusing on the reporting of reviews evaluating randomized trials, but also used as a basis for reporting systematic reviews of other types of research, particularly evaluations of interventions. For more information visit: www.prisma-statement.org/

Additionally, studies performed in Africa related to meningococcal A vaccine were excluded for not being relevant to Latin American and the Caribbean, as were those studies specifically targeting patients with sickle cell disease, HIV-infection, or conditions known to affect immune response. Others excluded were: studies that considered only the disease of selected serogroups without denominators, adverse events, or immunogenicity (antibody levels); studies of nasopharyngeal carriage; studies considering only laboratory data; studies with all-cause mortality and hospitalization as primary outcomes; and studies that assessed only nosocomial infections.

Study selection


Citations were screened by two independent reviewers in a two-step process. First, titles and abstracts were reviewed for duplication and inclusion/exclusion criteria. After eliminating duplicates, the full texts of these papers were obtained to complete the eligibility screening. Second, five citations on which eligibility reviewers disagreed were discussed or assessed by a third reviewer.

By screening the abstracts of the remaining 209 studies, 158 were excluded—30 were not related to meningococcal vaccine; 95 did not measure vaccine impact; 6 were reviews (not primary studies); 3 were reports or press release; 8 were studies of meningococcal A vaccine; 3 were studies of vaccines that are not commercially available; and 13 were modelling/economic evaluations.

The full texts of 51 studies were then evaluated, and another 19 were excluded, resulting in 32 primary studies for this review (12 – 43). Figure 1 shows the PRISMA Flow Diagram of the search strategy and selected articles.

Data collection and study quality

Data extraction was done independently by three reviewers, using abstraction forms developed specifically for this review. Data extracted included first author, country and state when available, vaccine and serogroup type, study type, data source, study period, vaccine introduction date, vaccination strategy, vaccination schedule used, age groups on which impact was measured, number of subjects on which impact was measured, case definition, effectiveness, or impact

FIGURE 1. PRISMA Flow Diagram for selection of articles for a review of the literature on the impact and/or effectiveness of meningococcal vaccines, 2016

Source: Prepared by the authors from study data.

as reported, and main conclusion of the study.

The main results included were impact and/or effectiveness assessments, corresponding confidence limits when available, and any information on control and ascertainment of exposure for case-control studies and cohort studies. Additionally, control (as steady practice) and presence of time series before were evaluated for impact studies.

Data analysis

A descriptive analysis was conducted of each study's characteristics including design, type, and schedule of meningococcal vaccine introduced, vaccination strategy, data sources, and endpoints considered. For all studies, the main measure of interest was the vaccine effect in reducing the outcome of interest.

RESULTS

Table 1 summarizes the characteristics of the 32 articles selected. Most of them were spread out evenly through the

study period. Of the total, 26 evaluated conjugate vaccines (12 – 37) and 6 evaluated polysaccharide vaccines (38 – 43). By far, the vaccine most commonly evaluated was the meningococcal conjugate C vaccine (Men-C-C), with 25 articles; some studies evaluated multiple vaccines. The studies originated from several countries, with Spain being the most prolific publisher (10 studies), followed by Canada (8 studies). The study periods of the 32 selected articles ranged from 1 – 58 years, with an average of 8.7 years and a median of 5 years. The most common data sources for the studies were surveillance systems (27 studies).

Regarding the immunization strategy, most countries studied had introduced a meningococcal vaccine to their routine immunization program (25 studies) and 4 had also conducted catch-up for older age groups. Eleven studies evaluated the use of a meningococcal vaccine in a campaign. The vaccination schedule used varied widely. In September 2015, the United Kingdom became the first country to introduce meningococcal vaccine (4CMenB), a protein sub-unit vaccine,

TABLE 1. Characteristics of the articles included in a review of the literature on the impact and/or effectiveness of meningococcal vaccines, 2016

Characteristics	n (Total 32)	%
Publication year		
1999	1	3
2000	1	3
2001	3	9
2002	0	0
2003	3	9
2004	4	13
2005	2	6
2006	0	0
2007	0	0
2008	1	3
2009	2	6
2010	0	0
2011	2	6
2012	2	6
2013	0	0
2014	5	16
2015	2	6
2016	4	13
Type of vaccine (not mutually exclusive)		
Meningo Conjugate C	25	78
Polysaccharide A+C	4	13
MenACWY-135	2	6
4CMenB	1	3
VA Meningo BC	1	3
Polysaccharide Menc	1	3
Country		
Spain	10	31
Canada	8	25
United Kingdom	4	13
Brazil	3	9
Italy	3	9
Australia	1	3
Cuba	1	3
Ireland	1	3
Scotland	1	3
Study type (not mutually exclusive)		
Impact	30	94
Effectiveness	14	47
Data sources		
Surveillance	27	84
Hospital records	5	16

using a 2- and 4 months-old schedule with an opportunistic catch-up at 3 and 4 months of age. Among the 26 studies that looked at the conjugate meningococcal vaccines (12 – 37), 23 evaluated routine introductions to national schedules and 6 evaluated campaigns; just a few studies evaluated both. Regarding the routine introduction of conjugate vaccines, the

most common schedule used was a single dose at 12–15 months of age (11 studies), followed by a three-dose schedule of 2-, 4-, and 6-months of age (6 studies). Four studies also vaccinated adolescents at around 12 years of age. Almost one-half of the conjugate vaccine schedules (12 articles) included a catch-up strategy for older age groups. Regarding the use of conjugate vaccines in campaigns, the targeted ages ranged from 2 months – 20 years of age.

The use of polysaccharides vaccines in the studies identified was predominantly during campaigns (4 of 6 studies), covering a wide range of age groups, usually including infancy – late adolescence. It is noteworthy that these studies were published in the 1990s (38–43).

Regarding the estimation of impact or effectiveness of the vaccines, approximately two-thirds of the studies (19 articles) evaluated the impact and/or effectiveness of the vaccine on the entire population, whereas the remaining one-third (13 articles) restricted the measure of effect to a certain age range or specific group. Meningococcal cases with lab confirmation (culture and/or cerebrospinal fluid polymerase chain reaction) were used as the most common endpoint included (25 studies) and secondary data from health information systems, surveillance systems, and other sources were also used. Seven studies evaluated laboratory-confirmed invasive meningococcal disease.

Fourteen studies measured effectiveness by comparing risk or rates in vaccinated and unvaccinated people during the same time period. Thirty studies measured impact by comparing a pre-vaccine introduction period to a post-vaccine introduction period. Regarding the degree of impact of meningococcal vaccines, one-third (11 studies) did not report a measure of change, but rather estimates before and after the introduction. Among the 21 studies that did report a specific measure of impact or change, the actual measure varied, with vaccine effectiveness being the most common measure (14 studies), followed by incidence decline (9 studies), and lastly, relative risk (2 studies). The age range on which the impact was measured also varied within and among studies. A summary of main outcomes and characteristics of the 32 studies included can be found in Tables 2a–d. Among the studies that estimated effectiveness of the polysaccharide vaccines, results ranged

from 65% (Canada)–83.7% (Spain). In the Canada study, vaccine effectiveness was not sustained beyond the second year of the study, showing no effectiveness in years 3–5. The only other study that provided an impact estimate for the polysaccharide vaccines was the United Kingdom, reporting a relative risk of 0.94.

With the studies that estimated the effect of the conjugate vaccines, effectiveness estimates ranged from 66% (United Kingdom)–100% (Spain), and the incidence decline ranged from 77% (Canada)–100% (Spain). The only study that evaluated the protein sub-unit vaccine reported a vaccine effectiveness of 82.9%.

DISCUSSION

Overall, the 32 studies included in this review showed a positive impact and effectiveness of both the polysaccharide vaccines and the conjugate vaccines on vaccine-serogroup meningococcal disease and mortality; but the conjugate vaccines show higher impact and effectiveness, plus longer lasting protection over the polysaccharide vaccines. Pelton and colleagues (3) reported similar results, including several studies that showed reductions in vaccine serogroup invasive meningococcal disease incidence after meningococcal polysaccharide vaccines or polysaccharide protein conjugate vaccine introduction, while the overall incidence of other serogroups remained stable, an effect observed in this review's studies as well. Nonetheless, it is important to consider the advantages of conjugate vaccines, which are well established in regards to immunologic memory and protection in the youngest children (less than 2 years of age) who respond poorly to polysaccharide vaccines. Also, the ability of conjugate vaccines to reduce carriage allows for significant indirect benefits through herd protection, reducing transmission of meningococci (44).

Study limitations

As mentioned above, this literature review selected effectiveness and impact studies of meningococcal vaccines. Vaccine effectiveness is a measure of protection attributable to a vaccine administered under field conditions to a given population and is measured by observational studies. In turn, impact of a vaccination program is measured by the fraction of

prevented disease in a population and is estimated by comparing populations with and without vaccination, usually the same population before and after vaccination. The impact of a vaccination program refers to the program's overall effect. Different study designs are used for measuring effectiveness and impact, but all the post-licensure study designs present a certain level of bias. For example, in effectiveness studies, the level of natural protection against disease may differ between vaccinees and non-vaccinees, the target group for vaccination may have a higher probability of developing the disease, or the ascertainment of the disease may differ depending on vaccination status. On the other hand, impact studies could enable bias, such as secular trends in baseline transmission or cyclical variations of the target disease, changes in case detection or surveillance methods, or the presence of a concomitant intervention, such as prudent antibiotic use, which has an impact on the target disease as well (45). The majority of the studies included in this review are impact studies using mostly before-and-after design. A study-by-study assessment of risk bias was not conducted and may represent a limitation to the interpretation of results.

Regarding the reported results, many of the studies did not report a measure of change; among those that did, their measures varied, as did the age range of their study participants. These variations make it difficult to draw comparisons between studies, and even harder to consolidate estimates in an attempt to obtain a "big picture" of the impact of meningococcal vaccines on disease and death due to meningococcus. Additionally, it is worth noting that the countries studied use different vaccine schedules, target a variety of age groups, implement primary vaccination with or without catch-ups, and apply diverse meningococcal disease diagnostic methods, all of which influence results and interpretation.

Of the 32 selected, there was only one study of the impact of the protein sub-unit vaccine serogroup B 4CMenB (35); this was in part because it was just licensed in 2015. The United Kingdom was the first country to introduce it to the national immunization program (46). During the program's first 10 months, the two-dose 4CMenB schedule has been highly effective in preventing meningococcal B (MenB) disease in infants.

TABLE 2a.

First Author, year	Country and state	Type of vaccine	Serogroups included in the vaccine	Specific vaccine	Study type	Data source	Study period	Vaccine introduction or campaign DATE
Lawrence, 2016	Australia	Conjugate	C	MCCV	Impact	Notification data	2000-2012	2003
Tauil, 2014	Brazil , Federal District	Conjugate	C	MCCV	Impact	Surveillance data; mortality database; public health laboratory database	2005-2011	2010
Cardoso, 2012	Brazil , Salvador	Conjugate	C	MCCV	Impact and effectiveness	Surveillance	Jan 2000- Dec 2011	1 Feb 2010
Bettinger, 2009	Canada	Conjugate	C	MCCV	Impact	Surveillance standardized case report forms from the national Notifiable Diseases Information System (SINAN)	2002-2006	Staggered implementation starting with campaigns in some provinces in 1989 to 2001, and routine programs starting in 2002 to 2005
De Wals, 2011	Canada	Conjugate	C	MCCV	Effectiveness	Laboratory and surveillance data	19 years	2001
Sadarangani, 2014	Canada	Conjugate	C	MCCV	Impact	Population based surveillance	1998-2012	2001 - recommendation in Canada
Sid, 2008	Canada , British Columbia	Conjugate	C	MCCV	Impact	IMD cases are reported to BC Centre for Disease Control Surveillance Denominator was obtained from an estimation of BC annual population using Population Extrapolation for Organization Planning	2003-2006	2003
Wormsbecker, 2015	Canada , Ontario	Conjugate	C and A C Y W	MCCV and ACWY	Impact	Surveillance data were obtained from the Ontario reportable diseases system, the integrated Public Health Information System (PHIS), and Public Health Ontario Laboratories (PHOL) for 2000-2013.	2000-2013	2004
Kihlin, 2009	Canada , Ontario, Quebec	Conjugate	C	MCCV	Impact	IMD cases reported to the central public health lab	2000-2006	After 2001
De Wals, 2004	Canada , Quebec	Conjugate	C	MCCV	Impact and effectiveness	Population-based observational study of cases of invasive serogroup C meningococcal	1996-2002	2001
O Maoldomhnaigh, 2016	Ireland	Conjugate	C	MCCV	Impact	Records from two tertiary pediatric hospitals	2001-2011	1 Oct 2000
De Waure, 2015	Italy	Conjugate	C	MCCV	Impact	Two surveillance sources: the informative System of Infectious Diseases (SIMI) since 1991, and the National Surveillance of Bacterial Meningitis since 1994	1994-2012	2005
Pascucci, 2014	Italy , Emilia Romagna	Conjugate	C	MCCV	Impact and effectiveness	Surveillance data	2000-2005	In Emilia-Romagna region: at risk - 2003; in 2006 routine to 12-15 mo and 14-15 yrs. No catch-up
Bechini, 2012	Italy , Tuscany	Conjugate	C	MCCV	Impact	Surveillance, The Regional Health Authority supplied all surveillance data on invasive bacterial disease (IBD) of the resident Tuscan population in the same age groups according to the Italian National Institute for statistics (ISTAT) data	2009-Nov 2011	1 Mar 2005
Mooney, 2004	Scotland	Conjugate	C	MCCV	Impact and effectiveness	Statistics Division of the Common Services Agency, National Health Service for Scotland Scottish Meningococcus and Pneumococcus Reference Laboratory,	1994-2003	1999

(Continued)

TABLE 2a. (Continued)

First Author, year	Country and state	Type of vaccine	Serogroups included in the vaccine	Specific vaccine	Study type	Data source	Vaccine introduction or campaign DATE
Cano, 2004	Spain	Conjugate	C	MCCV	Impact	Epidemiological surveillance of meningococcal disease in Spain is based on a passive notification system	1999/2000 to 2002/2003
Garrido, 2014	Spain	Conjugate	C	MCCV	Impact and effectiveness	Data on cases of meningococcal disease reported in Spain were obtained from the National Notifiable Disease Surveillance System Surveillance	January 2001 to December 2013
Larrauri, 2005	Spain	Conjugate	C	MCCV	Impact and effectiveness	Data on cases of meningococcal disease reported in Spain during the study period (1999–2004) were obtained from the Notifiable Disease Surveillance System	1999/2000 to 2001/2003
Salleras 2003 (Impact...)	Spain	Conjugate	C	MCCV	Impact	System population estimates computed at the midpoint of the periods considered and furnished by the National Statistics Institute (Instituto Nacional de Estadística)	Jan to Dec 2001
Cruz Rojo, 2005	Spain, Andalucia	Conjugate	C	MCCV	Impact	Andalucía meningococcal Surveillance and state demographic data	1997-2004
Salleras 2003 (Dramatic...)	Spain, Catalonia	Conjugate	C	MCCV	Impact and effectiveness	Surveillance data	1999–2000 To 2000–2001 And 2001–2002
Romero, 2011	Spain, Galicia	Conjugate	C	MCCV	Impact	Meningitis surveillance system	1990 - 1995
Morales, 2016	Spain, Navarra	Conjugate	C	MCCV	Impact and effectiveness	Surveillance data	1995-1999 and 2001-2014
Parikh, 2016	UK	Protein Sub Unit B Vaccine	B	4CMenB	Impact and Effectiveness	Surveillance	Sept 2015 to June 2016 1 Sep 2015
Martin, 2014	UK	Conjugate	C	MCCV	Impact	Routinely collected administrative statistics for hospital care	1963-1998 and 2000-2011
Trotter, 2004	UK	Conjugate	C	MCCV	Effectiveness	Cover of Vaccination Evaluated Rapidly (COVER) Child Health Department and National laboratories	1 Nov 1999

TABLE 2b.

First Author, year	Country and state	Specific vaccine	Vaccination STRATEGY (Routine or campaign?)	Vaccination schedule used	Age groups on which impact was measured	Case definition	Effectiveness or impact as reported	Impact estimation	Conclusion of the study (short)
Lawrence, 2016	Australia	MCCV	Routine	12 months and catch up for 2 to 19 year-olds	All ages	Invasive meningococcal disease and mortality	Incidence declined 96%, CI 94–98, and deaths declined from 68 in 2000–2002 to 3 deaths in 2010–2012	incidence decline 96%	Despite published evidence of waning antibody over time, an ongoing single dose of meningococcal C conjugate vaccine in the second year of life following widespread catch-up has resulted in near elimination of serogroup C disease in all age groups without evidence of vaccine failures in the first decade since introduction
Taiui, 2014	Brazil, Federal District	MCCV	Routine	3m - 2 years	All ages	Clinical manifestations consistent with MD who met at least one of the following criteria: isolation of <i>Neisseria meningitidis</i> from the cerebrospinal fluid (CSF) or blood; positive immunodiagnostic test for <i>N. meningitidis</i> antigen in the CSF or blood; presence of Gram-negative diplococcus in the CSF; or association or not of purpura fulminans with meningitis	The average annual incidence rate was 2.0/100 000 inhabitants/year (range: 1.3–2.5 per 100,000) from 2005 to 2009 and 1.8 and 0.8/100 000 inhabitants/year in 2010 and 2011. The average annual incidence rates in children under one year of age were 30.6, 13.3 and 13.1/100 000 inhabitants/year and in children with two years of age were 14.4, 10.8 and 2.7/100 000 inhabitants/year from 2005 to 2009 and in 2010 and 2011, respectively	Not expressed	In conclusion, the MCCV strategy implemented in Brazil proved highly effective and had a strong direct impact on the target population. However, incidence and case fatality rates of MD remain high with a wide gap in the risk of dis-ease between poor and affluent areas pointing to the need for periodic adjustments and reevaluations of the current strategy
Cardoso, 2012	Brazil, Salvador	MCCV	Routine + catch-up	2 and 4 months, with booster in the second year of life, and catch up in 2010 for people aged 10 to 24	All age groups	Lab confirmed meningococcal disease	VE 100%	Low coverage in the population targeted for mass vaccination may have limited impact on ongoing transmission of serogroup C meningococcal disease despite high vaccine effectiveness	
Bettinger, 2009	Canada	MCCV	first campaigns and afterwards routine.	Varies by province. Quebec and Alberta 2, 4 and 6 or 12 months. Other provinces: 2 and 12 months, or only 12 months. Most provinces did catch up to children (varying ages, 9 to 18 year-olds)	All age groups	Lab confirmed invasive meningococcal disease	Rates declined from 0.41 (0.28–0.60) in 2002 to 0.07 (0.02–0.16) in 2006	Not expressed	A substantial decrease in group C incidence occurred in provinces with early MenC immunization programs. Serogroup C incidence remained stable in provinces without MenC programs. We found no evidence of serogroup replacement

(Continued)

TABLE 2b. (Continued)

First Author, year	Country and state	Specific vaccine	Vaccination STRATEGY (Routine or campaign?)	Age groups on which impact was measured	Vaccination schedule used	Case definition	Effectiveness or impact as reported	Impact estimation	Conclusion of the study (short)
De Wals 2011	Canada	MCCV	campaign	2 months to 20 years	2m to 20 years	Laboratory confirmed (culture, PCR, CSF, polymerase 2001)	Effectiveness 87.4% (CI 75.4-94.2%)	VE 87.4%	Results support current Canadian recommendation to provide booster vaccination for adolescents
Sadarangani 2014	Canada	MCCV	Routine	Current schedule: <1 yr. (three doses); 12-23 mo and 12-24 yrs (1 dose)	all ages	Admission to hospital and identification of <i>Neisseria meningitidis</i> from a sterile site	Between 2002 and 2005, the incidence of IMD was 0.14 per 100 000 per year for serogroup C and 0.33 per 100 000 per year for all other serogroups combined. Between 2009 and 2012, the incidence decreased by 77% to 0.03 per 100 000 per year for serogroup C ($P < .0001$), with no significant change in non-C disease	Incidence decline 77%	MCCV dramatically reduced the incidence of serogroup C IMD in Canada through both direct and indirect effects. The observation that disease incidence decreased with different schedules suggests that the doses at 12 months (common to all provinces) and adolescence (7 of 8 provinces studied) were critical in achieving disease control
Siu, 2008	Canada, British Columbia	MCCV	Routine	Infants and adolescents	All ages	Serogroup-specific invasive meningococcal disease	Average annual incidence of serogroup-C IMD has declined from 0.32/100 000 in 2003 to 0.07/100 000 in 2005 in this age group with a significant downward trend ($p=0.05$)	Not expressed	There is a decreasing trend of pediatric serogroup C invasive meningococcal disease and an increase in median age of serogroup C IMD cases since 2003, most likely explained by protection from immunization
Wormsbecker, 2015	Canada, Ontario	MCCV and ACWY135	Routine	1 year-old, 12 year-olds	All ages	Invasive meningococcal disease	Reductions in serogroup C IMD among program eligible and ineligible age groups suggest both direct and indirect MCCV vaccine program impact	Incidence decline 17.2% per year	Reductions in serogroup C IMD over time (17.2% reduction per year [95% CI: 13.4 to 20.7]). The incidence of serogroup C IMD decreased significantly in children aged 1-17 years in the post-program period, based on age-specific incidence rate ratios (IRRs) and their 95% confidence intervals (CIs). Adolescents 12-16 years had the lowest serogroup C IRR (0.07/95% CI: 0.01 to 0.55); the rate decreased more than 14-fold between the pre- and post-periods.

(Continued)

TABLE 2b. (Continued)

First Author, year	Country and state	Specific vaccine	Vaccination STRATEGY (Routine or campaign?)	Vaccination schedule used	Age groups on which impact was measured	Case definition	Effectiveness or impact as reported	Impact estimation	Conclusion of the study (short)
Kinlin, 2009	Canada, Ontario	MCCV	Routine	12 months and catch up for children aged 12 or 15-19 year-olds	All ages	Invasive meningococcal disease	Decline from 5.48 to 4.26 cases per 1000 000. Rate of serogroup C decreased 50.12% while non-C decreased by 5.61%.	Rate of serogroup C decreased 50.12%	Conjugate serogroup C vaccination in Ontario appears to have had a direct effect on IMD incidence in children and adolescents, who are at greatest risk of invasive infection. The downward trend observed in older, unvaccinated age groups suggests that there are also herd benefits to immunization
De Wals, 2004	Canada, Quebec	MCCV	Campaign	One dose to all people aged 2 months to 20 years	All ages	Serogroup C meningococcal disease	The incidence rates were 1.04 per million in 1996-2000, 7.84 in 2001 (P=.001), and 3.63 in 2002. For the age group targeted for vaccination, the incidence increased from 2.90 per million in 1996-2000 to 21.47 in 2001, then decreased to 3.26 in 2002. For those 21 years and older, the incidence was similar between 2001 (3.26) and 2002 (3.77)	VE 96.8%	The new conjugate vaccine was effective in controlling an emerging epidemic of serogroup C meningococcal disease as well as providing short-term protection across a wide age range
O Maoldomhnaigh 2016	Ireland	MCCV	Routine	Under 19 years of age	Under 19 years of age	Menigococcal cases	From 14.75 per 100 000 (1999) to 2 per 100 000 (2011), and 98% of disease caused by serogroup B and a national CFR of 3.6%	Not expressed	Despite the meningococcal vaccination campaign, invasive meningococcal disease continues to cause serious morbidity and claim lives. Group B infections remain dominant. As children who die often present with fulminant disease, preventive strategies including use of meningococcal B vaccine are needed to avert death and sequelae
De Waure, 2015	Italy	MCCV	Routine + catch-up	13-15m; catch-up at 11-18yrs	<1 year; 1-4 yrs; 5-9 yrs; 10-14 yrs; 15-24 yrs; 25-64 yrs; 65 yrs and over	"Cases are defined according to clinical and laboratory criteria. A confirmed case is defined as a patient with a compatible clinical illness and a laboratory confirmation such as the detection -through microscopic direct examination, culture or polymerase chain reaction- of <i>N. meningitidis</i> from a normally sterile site or the identification of the polysaccharide antigen in the cerebrospinal fluid."	Not expressed	Our results suggest that MCC had an impact in decreasing the incidence of <i>N. meningitidis</i> C related IMD. However, data on typing are incomplete and efforts are needed to make them available for studying the need and the impact of other meningococcal disease	

(Continued)

TABLE 2b. (Continued)

First Author, year	Country and state	Specific vaccine	Vaccination STRATEGY (Routine or campaign?)	Age groups on which impact was measured	Vaccination schedule used	Case definition	Effectiveness or impact as reported	Impact estimation	Conclusion of the study (short)
Pascucci, 2014	Italy, Emilia Romagna	MCCV	2003 - at risk; 2006 - routine	<1yr; 1-4 yrs; 5-14 yrs; 15-24 yrs; 25-64 yrs; 65 and over; all ages	12-15 months and 14-15 years	Presence of N. meningitidis (either detected by standard culture or by DNA amplification of a pathogen specific genomic target) in a cerebrospinal fluid (CSF) sample with hospitalization	The average incidence of meningitis caused by meningococcus declined from 0.54/100 000 (years 2000–2005) to 0.33/100 000 (years 2006–2012). The total number of notified serogroup C cases dropped 0.2/100 000 in 2000–2005 to 0.06/100 000 (-70.1%) in 2006–2012. This change reached -100% (no case notified) and -83.1% in the two target age groups 1–4 and 15–24	Incidence related to serogroup C was 70.1%. VE was estimated as 90% or above in the age groups targeted to the vaccination campaign	No case of serogroup C related infection was observed since 2006 in children aged 1–4 years. These findings suggest that the single-dose vaccination strategy against serogroup C N. meningitidis targeted to the age groups 12–15 months and 14–15 years was effective in the Emilia-Romagna population. However, the occurrence of two cases of meningitis in a 5-month child and in a 9-years child suggests caution and careful consideration in surveillance for the next years
Becchini, 2012	Italy, Tuscany	MCCV	Routine	3 doses at 3, 5, and 13 months of age and catch up until age 6 with a single dose		Incidence rates were calculated for the following age groups: <1 year; 1–4 years; 5–14 years; 15–20 years; 21–30 years; 31–49 years; 50–64 years; ≥65 years.	invasive meningococcal C disease	Progressively increasing vaccination coverage	Not expressed
Mooney 2004	Scotland	MCCV	Routine	All people under 20 years of age		All ages	serogroup C meningitis disease	From 15.8 incidents per 100 000 subjects in 1999 (95% confidence interval [CI], 11.3–20.3) to 0.7 incidents per 100 000 subjects in 2001 (95% CI, 0.3 to 1.6), for subjects <5 years old, and from 6.7 incidents per 100 000 subjects in 1999 (95% CI, 5.1–8.3) to 1.5 incidents per 100 000 subjects in 2001 (95% CI, 0.7–2.3), for subjects 5–19 years old	VE 99.23%

(Continued)

TABLE 2b. (Continued)

First Author, year	Country and state	Specific vaccine	Vaccination STRATEGY (Routine or campaign?)	Vaccination schedule used	Age groups on which impact was measured	Case definition	Effectiveness or impact as reported	Impact estimation	Conclusion of the study (short)
Cano, 2004	Spain	MCCV	Routine	Infants (not further specified)	All Age groups	Lab confirmed meningococcal disease	RR 0.42	The incidence of meningococcal disease, especially serogroup C, has fallen sharply during the last three epidemiological seasons in Spain covered by this study	
Garrido, 2014	Spain	MCCV	Routine	2, 4, 6 months of age (in 2006, a booster dose during the second year was added)	All ages	Lab confirmed meningococcal disease	VE 94.83%	The meningococcal C conjugate vaccination programme has been extremely successful in controlling the disease and continues to be evaluated and adapted to the changes in the epidemiology of the disease to ensure long-term vaccine protection	
Larrauri, 2005	Spain	MCCV	Routine	2,4,6 months and catch up (variable ages)	Children under 6 years of age	Serogroup C meningococcal disease	incidence decline 85%. 97.8% VE for children vaccinated in the catch-up campaign and 95.2% for those who had been routinely vaccinated in infancy	The vaccine registered high short-term VE values but there has been some loss of VE with time. Four years after vaccination, vaccine protection levels exceeded 94% in cohorts immunised during the campaign. Among children vaccinated in routine childhood immunisation programmes, however, long-term VE loss was greater.	(Continued)

TABLE 2b. (Continued)

First Author, year (Impact...)	Country and state	Specific vaccine	Vaccination STRATEGY (Routine or campaign?)	Vaccination schedule used	Age groups on which impact was measured	Case definition	Effectiveness or impact as reported	Impact estimation	Conclusion of the study (short)
Salleras 2003 (Impact...)	Spain	MCCV	Routine and campaign	Routine: 2, 4, 6 months of age. Campaign: children aged between 2 months and 6 years	Children under 6 years of age	Lab confirmed meningococcal disease	Reported cases of serogroup C meningococcal disease decreased sharply after the mass vaccination campaign carried out during the year 2000 (18 cases in 2001 compared with 46 in 2000 observed in the <6 years age group (only 2 cases in 2001, both in non-vaccinated children), compared with 27 cases in 2000). Vaccination effectiveness in children <6 years was 100% (94.27–100%).	Not expressed	The results of this study show the high level of effectiveness of the meningococcal C conjugated vaccine in the short term
Cruz Rojo 2005	Spain, Andalucia	MCCV	Campaign	Under 10 years of age	<10 years	Meningococcal disease	The annual incidence in <10 years pre vaccination 8.2 per and post vaccination 2.0 per 100,000 inhabitants, this difference being statistically significant. Conversely, in the population over 10 years old has produced neither that downward trend in the incidence, nor a significant difference between the average annual rates of both periods (rates of 1.2 and 1.0 respectively).	Not expressed	The absence of vaccine failure and the impact observed on the incidence of serogroup C meningococcal disease in children under 10 suggests the effectiveness of this new conjugate vaccine
Salleras, 2003 (Dramatic...)	Spain, Catalonia	MCCV	Routine (2000) and campaign (2001, 2002)	Routine: 2, 4, 6 months. Campaign: 6-19 year-olds	All ages	Lab confirmed meningococcal disease	VE 100% The accumulated 4-weekly incidence rates in children <6 years of age show the dramatic decline in disease incidence during the 2000–2001 epidemic season a decline which was maintained during the 2001–2002 epidemic season. In contrast, in the 6–19 years age group, the moderate decrease observed during the 2000–2001 epidemic season was not maintained during the 2001–2002 epidemic season. Vaccination effectiveness in children <6 years was 100% (94.27–100%).	VE 100%	The available data seem to confirm that there has been no substitution of serogroup C by serogroup B after mass vaccination with the conjugated vaccine, at least in the short term. However, this subject still preoccupies expert opinion and only the future evolution of the disease will provide a definitive answer
Romero 2011	Spain, Galicia	MCCV	Routine infant; Catch-up	2,4,6 months; one time catch-up all children up to 19yrs; another one time catch-up 13–25 years who had not received the conjugate vaccine	13–25 yrs	Isolation of <i>Neisseria</i> meningitidis in a totally sterile environment, or detection of <i>Neisseria</i> meningitidis genome in a sterile environment or detection of <i>Neisseria</i> meningitidis in CSF or diplococcus gram-negative display in CSF	Incidence in 2004–2005 was 0.84 cases per 100 000 - reduced to 0.18 cases per 100 000 in 2007–2008	Not expressed	In Galicia, a series of vaccination campaigns, particularly focusing on high-risk groups, has shown high effectiveness, with a marked reduction in the disease incidence in the vaccination cohort accompanied by a relevant reduction in the overall population.

(Continued)

TABLE 2b. (Continued)

First Author, year	Country and state	Specific vaccine	Vaccination STRATEGY (Routine or campaign?)	Vaccination schedule used	Age groups on which impact was measured	Case definition	Effectiveness or impact as reported	Impact estimation	Conclusion of the study (short)
Morales, 2016	Spain, Navarra	MCCV	Routine + catch-up	Routine: newborns/ infants; catch-up campaigns: in 2000 - <6 yrs and in 2004 <17 yrs	<15 yrs and adults (all ages above 15yrs)	Presence of <i>N. meningitidis</i> via PCR, isolation from CSF, identification of <i>N. meningitidis</i> from a sterile site	Between 1995 and 1999, the mean annual incidence of meningococcus C disease was 20.81 and 14.13 cases per 100 000 in children younger than 1 year and 1 to 4 years, respectively. The first part of the post-vaccination period (2001-2003), the incidence was reduced by 100% in the 2 groups less than 5 years old ($p <0.001$). In the second part of the post-vaccination period (2004-2014), declines in the incidence of 95% were consolidated in the 1-year-olds ($p <0.001$) and 100% in the 1-to-4-year-olds ($P <0.001$)	Incidence decline 100% and 95%. VE 96%	The MenCC vaccination program has been successful in decreasing the incidence rate of serogroup C meningococcal disease in Navarra, and schedule changes have maintained high vaccine effectiveness throughout the study period
Parikh, 2016	UK	4CMenB	Routine	2, 4 months of age with opportunistic catch-up for 3 and 4-months old	Infants born on or after May 2015	Lab confirmed meningococcal disease	Two-dose vaccine effectiveness was 82.9% (95% CI 24.1-95.2) against all MenB cases	50% incidence reduction. VE 82.9%	The two-dose 4CMenB priming schedule was highly effective in preventing MenB disease in infants. Cases in vaccine-eligible infants halved in the first 10 months of the programme
Martin, 2014	UK	MCCV	Routine + catch-up	"Infant"; catch-up 19 yrs	<15 yr	"We defined diseases according to International Classification of Disease (ICD) codes (table 1) and included cases if the diagnosis was recorded as the primary or one of the secondary diagnoses."	Admission rates decreased by 66%.	Vaccine-preventable invasive bacterial disease in children has decreased substantially in England in the past five decades, most notably with the advent of effective conjugate vaccines since the 1990s. Ongoing disease surveillance and continued development and implementation of vaccines against additional pneumococcal serotypes and serogroup B meningococcal disease are important	
Trotter, 2004	UK	MCCV	Routine	2,3,4 months and catch-up to children younger than 18 years of age	All ages	Laboratory-confirmed meningococcal serogroup C disease	VE 66% to 83%	Vaccine effectiveness was high (83%) in all children who had received MCC vaccines in the catch-up campaign at age 5 months to 18 years. Overall, routine infant vaccination was estimated to be 66% effective, but clear differences were noted according to time since vaccination	

TABLE 2c.

First Author, year	Country and state	Type of vaccine	Serogroups included in the vaccine	Specific vaccine	Study type	Data source	Study period	Vaccine introduction or campaign date
Kupak, 2001	Brazil , Santa Catarina	Polysaccharide	C	MenC polysaccharide	Impact and effectiveness	National Surveillance system	March 1995 to March 1997	1 March 1996
De Wals, 2001	Canada , Quebec	Polysaccharide	A, C, Y, W	ACYW and A+C	Impact and effectiveness	Cases reported to regional health authorities by clinicians, hospital laboratories, and the provincial laboratory serving as a reference centre for <i>N. meningitidis</i>	1990-1998	Dec 1992 to March 1993
Perez-Rodriguez, 1999	Cuba	Polysaccharide vaccine	B+C	VA-Meningo-BC	Impact	Direct Information System	1991-1996	1991
Góicoechea Saez, 2003	Spain , Valencia	Polysaccharide vaccine	A+C	Menpox A+ C (polysaccharide)	Impact and effectiveness	Clinical records from public hospitals	1996-2000	Sep to Dec 1997
Pereiro, 2001	Spain , Valencia,	Polysaccharide	A+C	A+C (two different vaccines were used)	Impact	Population-based active surveillance	1996-1998	Oct to Dec 1997
Bergman, 2000	UK	Polysaccharide	A+C	Menpox A+ C (polysaccharide)	Impact	Hospital inpatient summaries	1989-1997	Late 1992

TABLE 2d.

First Author, year	Country and state	Specific vaccine	Vaccination STRATEGY (Routine or campaign?)	Vaccination schedule used	Age groups on which impact was measured	Case definition	Effectiveness or impact as reported	Impact estimation	Conclusion of the study (short)
Kupek, 2001	Brazil, Santa Catarina	MenC polysaccharide	Campaign	All children between 6 months and 14 years of age	All ages	Lab confirmed meningococcal disease	In general population of Santa Catarina, VE was 74.3% (52.7% to 99.6%), and in children 6 months to 14 years, VE was 93.1% (85.2% to 100%)	VE 74.3% and for children 6 months to 14 years of age 93.1%	Group C meningococcal vaccine is effective in reducing the occurrence of meningococcal disease in children 6 months to 14 years of age, and that the ratio of rate ratios (RRR) is a useful method to evaluate vaccine effectiveness
De Wals, 2001	Canada, Quebec	ACYW and A+C	Campaign	All persons between 6 months and 20 years of age	All ages	Meningococcal disease between 1990-1998 and culture-proven serogroup C meningococcal disease between 1993-1998	The incidence of serogroup C disease decreased after the mass immunization campaign, from 1.4 per 100 000 in 1990-1992 to 0.3 per 100 000 in 1993-1998, and the overall incidence of other serogroups remained stable at 0.7 per 100 000, with a small increase in the proportion of cases caused by serogroup Y. Protection from serogroup C MCD was indicated in the first 2 years after vaccine administration (VE, 65%; 95% confidence interval [CI], 20%-84%), but not in the next 3 years	VE, 65% first 2 years, afterwards 0%	Serogroups C polysaccharide vaccine is effective for controlling outbreaks in teenagers but should not be used in children younger than 2 years. The mass campaign did not induce significant serogroup switching
Perez-Rodriguez, 1999	Cuba	VA-Meningo-BC	Routine	3 and a half months, 5 and a half months	Children 1 to 4 years of age	Meningococcal cases with lab confirmation	Incidence density in 1991: 10.8 per 100 000. Incidence density in 1995: 0.67 per 100 000. Incidence density in 1996: 0.68 per 100 000.	Not expressed	The changes observed in the incidence of meningococcal disease and the displacement of higher risk to older age groups of 3 and 4 years of age constitute an important support to continue the administration of the vaccine in the routine program, even more so if a booster dose is added to the schedule
Goicoechea Saez, 2003	Spain, Valencia	Menpovax A+C (polysaccharide)	Campaign	18 months to 19 years of age	Children under 15 years of age	Invasive meningococcal disease	The rate of incidence by serogroup C in children under age 15 dropped following the vaccination campaign from 5.82/105 habitants in 1997 to 1.68/105 habitants in 1998. Rates similar to those prior to the time prior to the vaccination recorded three years subsequent to the campaign, showing an increase in the disease caused by serogroup B over the last 2 years. Sixty-one percent of the sequelae were among children under 5 years of age. Lethality was higher for serogroup C. Vaccination efficacy three years subsequent to the campaign was 83.7% for the 5-14 age range and 99.1% for the 19-month-4 year age range	VE 83.7% for 5-14 years and 69.1% for 19 m-4 years	The polysaccharide vaccine was shown to be effective for halting the outbreak. The drop in the incidence of serogroup C can be attributed to the vaccination efficacy achieved

(Continued)

TABLE 2d. (Continued)

First Author, year	Country and state	Specific vaccine	Vaccination STRATEGY (Routine or campaign?)	Vaccination schedule used	Age groups on which impact was measured	Case definition	Effectiveness or impact as reported	Impact estimation	Conclusion of the study (short)
Pereiro, 2001	Spain , Valencia,	A+C (two different vaccines were used)	Campaign	Persons between 18 months and 19 years of age living in the region	Children under 15 years of age	Meningococcal disease lab confirmed or clinical manifestations	The cumulative incidence increased from 10.6: 100 000 in 1996 to 15.2: 100 000 in 1997, but decreased to 7.9:100 000 in 1998, primarily due to a reduction in the incidence of serogroup C disease	Not expressed	Meningococcal polysaccharide vaccine seems to be an effective public health tool for the management of this serious communicable disease
Bergman, 2000	UK	Menpovax A+C (polysaccharide)	Routine	All recruits	1 dose of vaccine to all recruits	Bacterial meningitis or meningococcmia	The crude relative risk for meningococcal infection in recruits after vaccination compared with the period before vaccination was 0.94 (95% confidence interval (CI) 0.40 - 2.22) and the corresponding relative risk for trained personnel was 0.20 (95% CI 0.09 - 0.44)	RR 0.94	From 1993 onwards, no further clusters of group C infection were reported and incidence of meningococcal disease among trained soldiers fell, but there was no significant reduction in the overall incidence of meningococcal disease in recruits

Only four of the studies selected for this review originated in Latin America and the Caribbean (LAC). This was not unexpected since only three countries in the area have introduced a meningococcal vaccine into their routine schedule and conducting these types of studies in LAC is challenging due to the area's insufficient surveillance of meningococcal disease. All four studies obtained their disease-burden data from Regional public health laboratories. This reinforces the premise that regional surveillance systems are of paramount importance to estimating the impact of vaccine introduction, and therefore, should be adequately maintained and strengthened (13).

CONCLUSIONS

These 32 studies have suggested or shown a reduction in cases and deaths due to vaccine-serotype invasive meningococcal disease after introduction of a meningococcal vaccine to the routine schedule or as part of a mass campaign. When considering the meningococcal vaccine, each LAC country should take into account its particular epidemiologic profile, as well as the cost of these vaccines, only one of which—the conjugate tetravalent ACYW-135—is currently offered through the PAHO Revolving Fund (US\$ 20.30 per dose; 47).

LAC countries should also strengthen their epidemiological surveillance systems to include serogroup-specific surveillance of invasive meningococcal disease. Adequate surveillance can measure the magnitude of this public health issue, thereby informing decision makers, while setting an epidemiologic foundation for conducting future vaccine impact and effectiveness studies.

Conflict of interests: None declared.

Disclaimer. Authors hold sole responsibility for the views expressed in the manuscript, which may not necessarily reflect the opinion or policy of the RPSP/PAJPH and/or PAHO.

REFERENCES

- Plotkin S, Orenstein W, Offit P. *Vaccines*. 6th ed. Paris: Elsevier; 2012.
- Halperin SA, Bettinger JA, Greenwood B, Harrison LH, Jelfs J, Ladhami SN, et al. The changing and dynamic epidemiology of meningococcal disease. *Vaccine*. 2012;30 Suppl 2:B26–36.
- Pelton SI. The global evolution of meningococcal epidemiology following the introduction of meningococcal vaccines. *J Adolesc Health*. 2016;59(2 Suppl):S3–S11.
- Baccarini C, Ternouth A, Wieffer H, Vyse A. The changing epidemiology of meningococcal disease in North America, 1945–2010. *Hum Vaccin Immunother*. 2013;9(1):162–71.
- United States Food and Drug Administration. *Vaccines, Blood & Biologics—Bexsero*, 2016. Available from: <https://www.fda.gov/BiologicsBloodVaccines/Vaccines/ApprovedProducts/ucm431374.htm> Accessed 17 October 2017.
- European Medicines Agency. *Human medicines—Bexsero*, 2017. Available from: www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/002333/human_med_001614.jsp&mid=WC0b01ac058001d124 Accessed 17 October 2017.
- World Health Organization. WHO vaccine-preventable diseases: monitoring system. 2016 global summary, 2017. Available from: http://apps.who.int/immunization_monitoring/globalsummary Accessed 17 October 2017.
- World Health Organization. Immunization, vaccines and biologicals. Data, statistics and graphics. 2017; Available from: www.who.int/immunization/monitoring_surveillance/data/en/ Accessed 17 October 2017.
- Ruttmann RW, Gentile A, Parra MM, Saez-Llorens X, Safadi MA, Santolaya ME. A consensus statement: meningococcal disease among infants, children and adolescents in Latin America. *Pediatr Infect Dis J*. 2014;33(3):284–90.
- Safadi MA, Berezin EN, Oselka GW. A critical appraisal of the recommendations for the use of meningococcal conjugate vaccines. *J Pediatr (Rio J)*. 2012;88(3):195–202.
- Safadi MA, O’Ryan M, Valenzuela Bravo MT, Brandileone MC, Gorla MC, de Lemos AP, et al. The current situation of meningococcal disease in Latin America and updated Global Meningococcal Initiative (GMI) recommendations. *Vaccine*. 2015;33(48):6529–36.
- Lawrence GL, Wang H, Lahra M, Booy R, Mc IP. Meningococcal disease epidemiology in Australia 10 years after implementation of a national conjugate meningococcal C immunization programme. *Epidemiol Infect*. 2016;144(11):2382–91.
- Tauil Mde C, Carvalho CS, Vieira AC, Waldman EA. Meningococcal disease before and after the introduction of meningococcal serogroup C conjugate vaccine. Federal District, Brazil. *Braz J Infect Dis*. 2014;18(4):379–86.
- Cardoso CW, Pinto LL, Reis MG, Flannery B, Reis JN. Impact of vaccination during an epidemic of serogroup C meningococcal disease in Salvador, Brazil. *Vaccine*. 2012;30(37):5541–6.
- Bettinger JA, Scheifele DW, Le Saux N, Halperin SA, Vaudry W, Tsang R. The impact of childhood meningococcal serogroup C conjugate vaccine programs in Canada. *Pediatr Infect Dis J*. 2009;28(3):220–4.
- De Wals P, Deceuninck G, Lefebvre B, Boulianne N, De Serres G. Effectiveness of serogroup C meningococcal conjugate vaccine: a 7-year follow-up in Quebec, Canada. *Pediatr Infect Dis J*. 2011;30(7):566–9.
- Sadarangani M, Scheifele DW, Halperin SA, Vaudry W, Le Saux N, Tsang R, et al. The impact of the meningococcal serogroup C conjugate vaccine in Canada between 2002 and 2012. *Clin Infect Dis*. 2014;59(9):1208–15.
- Siu T, Tang W, Dawar M, Patrick DM. Impact of routine immunization using meningococcal C conjugate vaccine on invasive meningococcal disease in British Columbia. *Can J Public Health*. 2008;99(5):380–2.
- Wormsbecker AE, Wong K, Jamieson FB, Crowcroft NS, Deeks SL. Epidemiology of serogroup C and Y invasive meningococcal disease (IMD) in Ontario, 2000–2013: Vaccine program impact assessment. *Vaccine*. 2015;33(42):5678–83.
- Kinlin LM, Jamieson F, Brown EM, Brown S, Rawte P, Dolman S, et al. Rapid identification of herd effects with the introduction of serogroup C meningococcal conjugate vaccine in Ontario, Canada, 2000–2006. *Vaccine*. 2009;27(11):1735–40.
- De Wals P, Deceuninck G, Boulianne N, De Serres G. Effectiveness of a mass immunization campaign using serogroup C meningococcal conjugate vaccine. *JAMA*. 2004;292(20):2491–4.
- Ó Maoldomhnaigh C, Drew RJ, Gavin P, Cafferkey M, Butler KM. Invasive meningococcal disease in children in Ireland, 2001–2011. *Arch Dis Child*. 2016;101(12):1125–9.
- de Waure C, Miglietta A, Nedovic D, Mereu G, Ricciardi W. Reduction in Neisseria meningitidis infection in Italy after Meningococcal C conjugate vaccine introduction: A time trend analysis of 1994–2012 series. *Hum Vaccin Immunother*. 2016;12(2):467–73.
- Pascucci MG, Di Gregori V, Frasca G, Rucci P, Finarelli AC, Moschella L, et al. Impact of meningococcal C conjugate vaccination campaign in Emilia-Romagna, Italy. *Hum Vaccin Immunother*. 2014;10(3):671–6.
- Bechini A, Levi M, Boccalini S, Tiscione E, Balocchini E, Canessa C, et al. Impact on disease incidence of a routine universal and catch-up vaccination strategy against Neisseria meningitidis C in Tuscany, Italy. *Vaccine*. 2012;30(45):6396–401.
- Mooney JD, Christie P, Robertson C, Clarke SC. The impact of meningococcal

- serogroup C conjugate vaccine in Scotland. *Clin Infect Dis.* 2004;39(3):349–56.
27. Cano R, Larrauri A, Mateo S, Alcala B, Salcedo C, Vazquez JA. Impact of the meningococcal C conjugate vaccine in Spain: an epidemiological and microbiological decision. *Euro Surveill.* 2004;9(7):11–5.
28. Garrido-Estepa M, Leon-Gomez I, Herrero R, Cano R. Changes in meningococcal C epidemiology and vaccine effectiveness after vaccine introduction and schedule modification. *Vaccine.* 2014;32(22):2604–9.
29. Larrauri A, Cano R, Garcia M, Mateo S. Impact and effectiveness of meningococcal C conjugate vaccine following its introduction in Spain. *Vaccine.* 2005;23(32):4097–100.
30. Salleras L, Dominguez A, Cardenosa N. Impact of mass vaccination with polysaccharide conjugate vaccine against serogroup C meningococcal disease in Spain. *Vaccine.* 2003;21(7–8):725–8.
31. Cruz Rojo C, Garcia Gil C, Nieto Vera J, Monroy Morcillo A. Impact of systematic vaccination with the antimeningococcal C conjugated vaccine in a health area in Andalusia. *Rev Esp Salud Publica.* 2005;79(6):655–63.
32. Salleras L, Dominguez A, Cardenosa N. Dramatic decline of serogroup C meningococcal disease in Catalonia (Spain) after a mass vaccination campaign with meningococcal C conjugated vaccine. *Vaccine.* 2003;21(7–8):729–33.
33. Rego Romero E, Nartallo Penas V, Taboada Rodriguez JA, Malvar Pintos A, Hervada Vidal X, Lopez Pimentel MJ. Implementation and impact of a meningococcal C conjugate vaccination program in 13- to 25-year-old individuals in Galicia, Spain. *Z Gesundh Wiss.* 2011;19(5):409–15.
34. Morales D, Garcia-Cenoz M, Moreno L, Bernaola E, Barricarte A, Castilla J. Menin-gococcal C conjugate vaccine: Impact of a vaccination program and long-term effectiveness in Navarra, Spain, 2000–2014. *Enferm Infect Microbiol Clin.* 2016;34(10):639–44.
35. Parikh SR, Andrews NJ, Beebejaun K, Campbell H, Ribeiro S, Ward C, et al. Effectiveness and impact of a reduced infant schedule of 4CMenB vaccine against group B meningococcal disease in England: a national observational cohort study. *Lancet.* 2016;388(10061):2775–82.
36. Martin NG, Sadarangani M, Pollard AJ, Goldacre MJ. Hospital admission rates for meningitis and septicaemia caused by *Haemophilus influenzae*, *Neisseria meningitidis*, and *Streptococcus pneumoniae* in children in England over five decades: a population-based observational study. *Lancet Infect Dis.* 2014;14(5):397–405.
37. Trotter CL, Andrews NJ, Kaczmarski EB, Miller E, Ramsay ME. Effectiveness of meningococcal serogroup C conjugate vaccine 4 years after introduction. *Lancet.* 2004;364(9431):365–7.
38. Kupek E, Puricelli RC, Westrupp MH. Effectiveness of a mass immunization campaign against serogroup C meningococci in children in the federal state of Santa Catarina, Brazil. *Braz J Infect Dis.* 2001;5(6):324–31.
39. De Wals P, De Serres G, Niyonsenga T. Effectiveness of a mass immunization campaign against serogroup C meningococcal disease in Quebec. *JAMA.* 2001;285(2):177–81.
40. Perez Rodriguez A, Meneses FO. VA-MENGOC-BC vaccine: its impact on meningococcal disease in children 1–4 years old. *Rev Cubana Med Trop.* 1999;51(3):189–93.
41. Goicoechea Saez M, Fullana Montoro AM, Momparler Carrasco P, Redondo Gallego MJ, Brines Solanes J, Bueno Canigral FJ. Evolution of meningococcal infection among infant population in the autonomous community of Valencia (1996–2000). Effectiveness of A+C meningococcal vaccination. *Rev Esp Salud Publica.* 2003;77(1):125–42.
42. Pereiro I, Diez-Domingo J, Morant A, Gimeno C, Lerma M, San-Martin M, et al. Epidemiological pattern of meningococcal disease in Valencia, Spain. Impact of a mass immunization campaign with meningococcal C polysaccharide vaccine. *Scand J Infect Dis.* 2001;33(8):581–4.
43. Bergman BP, Hayton JC, Green AD. Effectiveness of the meningococcal vaccination programme for British Armed Forces recruits. *Commun Dis Public Health.* 2000;3(4):298–9.
44. Khatami A, Pollard AJ. The epidemiology of meningococcal disease and the impact of vaccines. *Expert Rev Vaccines.* 2010;9(3):285–98.
45. Hanquet G, Valenciano M, Simondon F, Moren A. Vaccine effects and impact of vaccination programmes in post-licensure studies. *Vaccine.* 2013;31(48):5634–42.
46. GOV.UK. Correspondence. MenB vaccination: introduction from September 2015. Available from: <https://www.gov.uk/government/publications/menb-vaccination-introduction-from-1-september-2015> Accessed 17 October 2017.
47. Pan American Health Organization. Revolving Fund Prices, 2017. Available from: http://www2.paho.org/hq/index.php?option=com_content&view=article&id=9561%3A2014-revolving-fund-prices&catid=839%3Arevolving-fund&Itemid=40714&lang=en Accessed 17 October 2017.

Manuscript received on 2 May 2017. Accepted for publication on 11 August 2017.

RESUMEN**Examen del impacto y la efectividad de las vacunas antimeningocócicas**

Objetivos. Resumir y evaluar críticamente los datos científicos sobre la repercusión y la eficacia de los programas de vacunación antimeningocócica en todo el mundo para orientar la toma de decisiones en América Latina y el Caribe.

Métodos. Se realizó un examen de la bibliografía siguiendo varios componentes de las directrices correspondientes a los elementos de notificación preferidos para revisiones sistemáticas y metanálisis. Se hicieron búsquedas en PubMed Central® para encontrar documentos publicados en cualquier idioma desde enero de 1999 hasta marzo del 2017.

Resultados. En total, se incluyeron 32 estudios, en cuya mayoría se evaluaba la vacuna antimeningocócica conjugada contra el serotipo C. En 14 estudios se midió la efectividad y en 30, el impacto. La efectividad de las vacunas polisacáridas se encontraba entre 65% y 83,7% (grupos etarios diferentes), mientras que la de las vacunas conjugadas, entre 66% y 100%. Gracias a la vacuna conjugada, la disminución de la incidencia de la enfermedad meningocócica confirmada en laboratorio varió entre 77% y 100% en diferentes grupos etarios. En el único estudio en el que se evaluó la vacuna de subunidades proteínicas se notificó que su efectividad era de 82,9%.

Conclusiones. Los estudios examinados muestran el impacto y la efectividad tanto de las vacunas polisacáridas como de las vacunas conjugadas en relación con la enfermedad meningocócica causada por los serotipos utilizados en las vacunas. Sin embargo, las vacunas conjugadas demuestran que tienen un mayor impacto y efectividad que las vacunas polisacáridas para dar una protección más duradera. En vista de la varianza del posible uso de una vacuna antimeningocócica, se deben fortalecer los sistemas de vigilancia epidemiológica para orientar las decisiones nacionales.

Palabras clave

Vacunas meningococicas; programas de inmunización; toma de decisiones, prevención & control; revisión; América Latina.

RESUMO**Impacto e efetividade das vacinas meningocócicas: uma revisão sistemática**

Objetivos. Sumarizar e analisar de maneira crítica as evidências sobre o impacto e a efetividade dos programas de vacinação contra doença meningocócica em todo o mundo para subsidiar a tomada de decisão na América Latina e no Caribe.

Métodos. Uma revisão da literatura científica foi realizada de acordo com vários componentes das diretrizes dos Principais itens para relatar revisões sistemáticas e meta-análises (PRISMA). Foi feita uma pesquisa da base de dados PubMed Central® em busca de estudos publicados em qualquer idioma de janeiro de 1999 a março de 2017.

Resultados. Trinta e dois estudos foram selecionados, sendo que a maioria avaliou a vacina conjugada contra o meningococo C. Quatorze artigos avaliaram a efetividade da vacina e 30, o impacto. A efetividade das vacinas polissacáridicas foi de 65% a 83,7% (em diferentes faixas etárias) e a das vacinas conjugadas foi de 66% a 100%. A redução com o uso da vacina conjugada da incidência de doença meningocócica confirmada por laboratório variou de 77% a 100% em diferentes faixas etárias. O único estudo que avaliou a vacina de subunidade proteica informou uma efetividade de 82,9%.

Conclusões. Os estudos examinados indicam impacto e efetividade da vacina polissacáridica e da vacina conjugada para doença meningocócica do respectivo sorogrupo vacinal. As vacinas conjugadas demonstram maior impacto e efetividade e proteção mais duradoura em relação às vacinas polissacáridicas. Diante da variação do uso em potencial da vacina meningocócica, os sistemas de vigilância epidemiológica devem ser reforçados visando subsidiar a tomada de decisão das autoridades nacionais.

Palavras-chave

Vacinas meningocócicas; programas de imunização; tomada de decisões, prevenção & controle; revisão; América Latina.