



ANNUAL RISK OF TUBERCULOUS INFECTION\*

by

G.M. Cauthen<sup>1</sup>, A. Pio<sup>2</sup> and H.G. ten Dam<sup>3</sup>

I. INTRODUCTION

Risk of infection in a population is in many respects the most informative index of the magnitude of the tuberculosis problem (Sutherland, 1976; Pio, 1984). The risk of infection at a particular time indicates the current magnitude of the incidence and prevalence of infectious cases (Styblo, 1985) and also indicates the magnitude of the tuberculosis problem years into the future. An observed decline in the risk of infection would be the earliest indicator of a decline in the epidemic cycle of tuberculosis, resulting from tuberculosis control activities or improvements in living standards. A rising risk of infection would be an early indicator of changes in the other direction, signalling the introduction of new risk factors, such as the spread of human immunodeficiency virus (HIV) infection.

It is known that risk of infection has been declining for many years in developed countries, but remains at high levels in many developing countries (Styblo, 1984). In developing countries in the last decade, a number of surveys of infection prevalence have been carried out in national populations as well as in smaller populations.

Therefore a project was carried out to assess the current level and trend in the risk of infection in developing countries by reviewing and assembling tuberculin skin test survey data available since 1975.

II. METHODS

Tuberculin skin test data collected since 1975 for populations in developing countries were assembled from reports to the World Health Organization and from the published literature. Data from surveys of childhood age groups that were judged of sufficient quality were selected in order to provide as valid and up to date an assessment as practical of the magnitude of the risk of infection. Comparable prior data for the same countries were also selected in order to judge whether the risk of infection is likely to have declined.

Prevalence of infection observed in childhood age groups was used to derive the average annual risk of infection that would have resulted in the observed cumulative prevalence rate. Choosing younger age groups allows the calculated average annual risk to be bracketed within a relatively narrow period of time between the average birth date of the group and the date of the survey (Styblo et al, 1969).

Each survey was judged on the basis of available documentation for ability to represent the population and to detect the proportion infected at a particular time. A sample survey was judged to have measured the proportion infected in the target population if the probability sampling design and the estimation method appeared correct in concept and conduct, and skin testing technique appeared adequate to measure the proportion infected in the sampling units.

1

Division of Tuberculosis Control, U.S. Centers for Disease Control, Atlanta, Georgia, USA.

2

Formerly Chief, Tuberculosis and Respiratory Infections Unit, Division of Communicable Diseases, World Health Organization, Geneva, Switzerland.

3

Scientist, Tuberculosis Unit, Division of Communicable Diseases, World Health Organization, Geneva, Switzerland.

\* The figures have been redrawn for reproduction by Dr Hans L. Rieder, Switzerland.

Basis for selection

Surveys were selected which met most of the following criteria:

Specified the sampling design including the sampling frame, staging, stratification, sampling units, allocation, sampling weights, and estimation formulas and documentation to support that the design was followed with adequate coverage.

Provided reaction-size distributions for surveyed age groups and for bacteriologically confirmed cases from the same population in order to assess the definition of infection and in order to judge technique.

Documented type and strength of antigen used, technique of administration and reading, and quality control procedures.

Specified the methods used to detect and eliminate persons with a history of BCG vaccination, and described BCG policy and actual coverage in the population age group surveyed.

Data assembled from selected surveys:

The area population represented and the survey period.

In the youngest age ranges for which sufficient numbers were tested and read, the number tested and read, the number counted as infected with *M. tuberculosis*, and the mean age.

Information documenting quality of the data.

Criterion of infection

Induration reaction size distributions observed in each survey were used together with general knowledge about the specific and nonspecific components of observable distributions to set a uniform criterion for counting infected persons in order to improve comparability between surveys.

Because it can be assumed that nonspecific reactions to intermediate strength purified protein derivative tuberculins are only infrequently larger than the mode of the distribution of true reactions, an observed distribution of tuberculin reactions will be nearly free of contamination by nonspecific reactions above its mode.

Because of this, and because distributions among truly infected persons are symmetric, the number truly infected in a distribution contaminated by nonspecific reactions would be closely approximated by adding the number at the mode plus twice the number with reactions larger than the mode.

This method, described by Nyboe (1960) and by Comstock et al. (1971), was used to determine the number of infected persons in the selected groups when detailed reaction size distributions were available and when the mode of reactions among infected persons in the same general population could be estimated from distributions of either bacteriologically confirmed cases or older age groups in the population.

When such data were not available, a cutoff criterion was accepted as the second choice if it appeared reasonably adequate based on distribution data.

Prevalence

Prevalence at a particular calendar time for an age group was calculated as the number infected divided by the number tested and read. Reported prevalence figures were used when numerator or denominator counts were not specified.

Risk of infection

For comparability, a single uniform method was chosen to derive the approximate average annual risk of infection (Styblo et al, 1969, pg. 14). By this method, the annual risk of infection (R) for a group of average age (A) was derived from the prevalence (P) by

$$R = 1 - (1 - P)^{1/A}.$$

The slope per year (B) of the trend between two risk estimates  $R_i$  and  $R_j$  at years  $Y_i$  and  $Y_j$  was approximated by

$$B = 1 - (R_j/R_i)^{1/T} \text{ where } T = Y_j - Y_i.$$

Average age

Average age was calculated by taking the midpoint of a single year (e.g., 6.5 for age given as 6 at last birthday) or a range of years (e.g., 7.0 for ages given as 6 to 7, or 7.5 for ages given as 5 to 9). When available, the weighted average age was calculated from year-of-age specific frequencies for an age group.

Time at which average risk occurred

For purposes of calculating and plotting trends, the calendar time of occurrence of a particular average annual risk ( $Y_R$ ) was considered to center approximately at the calendar time of the midpoint of the average lives of the individuals in the age-group. This was calculated by  $Y_R = Y_S - A/2$ , where  $A$  is the average age of the group and  $Y_S$  is the midpoint time of the survey. For example, a group of average age 6.1 years surveyed at 1978.9 was taken to contribute information about the annual risk of infection for a time interval centered at  $1978.9 - 6.1/2 = 1975.85$ .

The midpoint time of the survey ( $Y_S$ ) was calculated from the month, if stated, or the calendar year of the survey. For example, a survey stated to have taken place in 1975 was counted as occurring at 1975.5. A survey stated to have taken place from 1981 to 1982 was counted as occurring at 1982.0. A survey stated to have taken place from May 1952 to September 1953 was counted as occurring at 1953.04 (the midpoint between  $4.5/12 + 1952$  and  $8.5/12 + 1953$ ).

## III. RESULTS


A variety of tuberculin skin-test surveys were found in the published and unpublished literature: National sample surveys, sample surveys of other large populations, school surveys, school BCG campaigns, and mass BCG campaigns.

Twenty-five countries were judged for the purposes of this project to have been adequately surveyed in whole or significant part since 1975.

Figures 1 and 2 display the results for each country by region. Table 1 contains the data used in the figures and describes the area population represented, the survey design, the level of BCG coverage in the age-group surveyed, and the antigen and criterion defining infection.

An arithmetic scale is used in Figure 1 in order to display the relative magnitudes of current estimates of annual risk of infection.

Fig. 1. Annual risk of tuberculous infection in low-income countries of five WHO regions



A logarithmic scale of annual risks is used in Figure 2 in order to represent constant percentage decreases or increases between current estimates and prior estimates as straight line trends.

Dark symbols in the figures indicate risk estimates based on 42 well-conducted national sample surveys in 16 countries. In Figure 2, dark solid lines are used to connect rates which are based on more than one large sample survey of the same country.

Light symbols in Figure 2 indicate risk estimates based on data representative of lesser populations.

Dark dashed lines in Figure 2 are used to connect risk estimates based on national sample survey data with risk estimates based on other substantial data such as tuberculin testing in the mass BCG campaigns. A light solid line was used to connect risk estimates from repeated surveys representative of lesser populations. A light dashed line was used to connect surveys of approximately the same population done in different years.

The symbol + was used in Figure 2 to indicate the average of a set of risk estimates which together would represent a larger population. Stratum estimates within sample surveys, plotted as separate points to reveal variation in risk, were displayed as a single estimate for the whole population in this way. Sets of small area estimates that were less representative of the whole population than would have been provided by formal probability sampling, such as the set of completed sampling units in a not yet completed national survey, or areas selected by judgement from geographic or economic strata in order to be representative of a large heterogeneous population, were also combined in this way.

To indicate the extent of underlying variability, risk estimates from single surveys that were representative of only small areas are indicated in Figure 2 by unconnected light symbols.

#### Results by region

Africa. The most recent data for the eight countries included indicate current annual risks of infection around 1 to 2 percent. In the five countries for which comparable earlier surveys were available, trends over 20 to 25 years appear to be downward at 1 to 6 percent a year.

Data for the nearly completed national survey of Tanzania indicate an annual risk of infection of 1.1 percent, based on a population weighted average pooling of the samples. However, comparisons with earlier data available for three of the 18 regions in the sample do not together suggest a downward trend.

Surveys in Botswana and Lesotho suggest downward trends of about 6 and 1 percent a year, respectively. In the most recent survey in Botswana, however, BCG coverage in the age group surveyed had reached 83 percent, leaving only 17 percent who had by whatever circumstance missed vaccination, upon which to base the estimate.

No national surveys were available for Cameroon, but two surveys of the capital city, Yaounde, were available. These surveys indicate a level of annual risk of 0.6 percent in 1980 after a 20-year declining trend of 3.6 percent a year.

Survey data collected 30 years apart were available for Addis Ababa, the capital city of Ethiopia. These surveys indicate a 1.3 percent level of annual risk in 1979 succeeding a downward trend of 3.7 percent a year from the 4.1 percent risk level in 1949. A survey of selected villages in the Southwest, however, indicated a 3.8 percent annual infection risk in 1973. This suggests that within Ethiopia there is large variability in the level and perhaps also in the trend of infection risk.

Data from the 1976 sample survey of Gambia indicates an annual risk in 1971 of 1.9 percent. The trend of risk could not be assessed in Gambia, however. The only prior data available are from the 1958-59 survey of the capital city, Bathurst. The annual risk of infection in the urban stratum (4.4 percent) was higher than in the rural stratum (1.7 percent), which was nearly the same as the national level in 1971.

The annual risk of infection derived from skin test data for all military recruits in Burundi in 1981-82 was 1.2 percent. Data from an earlier survey among a scattered provincial population were selected in order to provide a rough comparison. The annual risk of infection in that group was 2.7 percent in 1955.

No comprehensive survey of the Nigerian population was available, but data from a survey of one emirate indicates an annual risk of 2.3 percent in 1973.



Americas. National sample survey data were available only from Argentina, but repeated surveys of large local populations within Argentina and also within Brazil were available. In the latest surveys, infection risks in these local populations varied ten-fold from 0.2 to 2 percent per year. Over time in these local populations, there was evidence of a steep decline in risk in Argentina, and a lesser decline in Brazil.

The annual risk of infection derived from the 1974-78 survey of the entire population of Argentina was 0.6 percent. Data from three surveys repeated in Santa Fe City and three surveys repeated in areas around Santa Fe City indicate low recent levels in that region of 0.2 to 0.3 percent annual risk, and indicate that the trend of the previous 12 to 15 years was steeply downward at a rate of 7 to 10 percent a year. The two trends described by the two sets of three survey points suggest that the decline was much steeper between the first two surveys but that the trend was nearly level during the 5 years between the second and third surveys. In the 1960-61 sample survey of another area of Argentina, Resistencia Province, the annual risk of infection was 0.5 percent.

Capitals of Brazilian States and Territories were surveyed in 1970-73 and again in 1980. The annual risk of infection was 0.8 to 1.9 percent in 1976 with evidence of a yearly downward trend of around 2 to 3 percent. A 1983 survey of the provincial population of Rio Grande Do Sul indicated a 0.4 percent annual risk of infection, with a 2.4 fold higher risk in southern than in northern rural areas.

South-East Asia. Only Thailand had national survey data available, but repeated surveys of local area populations were available for both India and Indonesia. In these populations, annual risks in the decade of the 1970's appear to have ranged from about 1 to 5 percent. Annual risks were highest in Indonesia.

Based on the 1977-79 national survey, the annual risk of infection in Thailand was 2.2 percent in 1974. Two earlier surveys, one of Bangkok and Chiangmai Province in 1960-64, and one of localities of Chiangmai and Kanchanaburi Provinces in 1954, indicate annual risks only slightly higher than in the national survey 16 to 24 years later.

In India, three large local populations were twice surveyed. Declining trends of 1.6 and 3.5 percent a year were apparent in two of these populations, but an increase of about 1 percent was apparent in the third population. Village surveys in 7 regions throughout India in 1972 indicate annual risks on the order of 1 to 2 percent. The village survey in Kashmir, compared to a 1978 sample survey of the Kashmir Valley, suggest a declining annual trend, but the populations may be only nominally comparable.

Indonesia has followed a policy since 1974 of withholding BCG and conducting 5-yearly surveys in nine special areas selected to be situated throughout the country. Three surveys have been completed in five of these areas, two in three areas, and one in another area. There are also three surveys available from an area with low BCG coverage. The latest surveys in these areas indicate a median annual risk of 2.3 percent (range: 0.7 to 3.9). Earlier data from the 1964-65 survey in rural East Java indicate an annual risk of 1.6 percent. Comparing the earliest to the latest survey in each of the nine areas with repeated surveys, the median downward trend in the annual risk of infection was 2.5 percent a year (ranging from a decline of 6.3 percent to an apparent increase of 3.4 percent a year).

Eastern Mediterranean (and Algeria). For the purposes of this project, Algeria is grouped in this region instead of with the African countries. National survey data were available for six countries, and repeated surveys of local populations were available for a seventh country in this region. Recent annual risk levels appear to be low to very low in Bahrain, Libya, Algeria and Kuwait (0.2 to 0.5 percent), but high in Afghanistan and Pakistan (1.8 and 3.5 percent, respectively). Recent levels of risk in the Syrian cities of Aleppo and Homs also appear to be very low (0.3 and 0.14, respectively). The trend of risk also appears to have decreased steeply in the current low risk countries, but not as steeply in Pakistan and appears to have increased in Afghanistan.

Four local areas sampled in 1980-84 for the nearly completed national survey of Algeria had also been canvassed in the 1949-52 mass BCG campaign. Repeated surveys were also available for the population of Blida. Based on these comparisons, annual risk of infection appears to have decreased about 6 to 7 percent a year in Algeria. A similar trend is apparent between risk estimates for the Syrian cities of Aleppo and Homs.

In Kuwait between 1972 and 1981, all school entrants (including non-citizens) were tuberculin tested prior to BCG vaccination. Compared to the 1962-63 national sample survey estimate, the annual risk of infection appears to have declined by 10 percent a year. In Bahrain, a decline of nearly 12 percent a year appears to have occurred between the 1969 national sample survey and mass tuberculin testing of students in 1981. A similarly steep rate of decline may have occurred in Eastern Libya, if the areas and populations covered in the surveys are comparable.

The annual risk estimate based on the 1974-78 national sample survey of Pakistan, which was completed except for Baluchistan, compared to the estimate based on the mass BCG campaign of 1949-54, suggests that the risk of infection declined by 4.2 percent a year, but that the risk was still high (1.8 percent) in 1972. The risk estimate based on the partially completed 1961-62 national survey is consistent with this trend.

The annual risk of infection in Afghanistan appears high (3.5 percent in 1978), without having much changed since 1958, compared with an estimate based on tuberculin testing program data collected in 1963 in selected schools around the country.

Western Pacific. Eleven national sample surveys were available for five countries in this region. In the Republic of Korea, the annual risk of infection appears to be declining, but remains high at about 2 percent. The risk appears to be about 1.8 percent in the Philippines, based on the recent national survey. Infection risk appears to be relatively low in Malaysia (0.4 percent) and Samoa (0.5 percent).

In China the annual risk of infection appears to have been 1 percent in 1975, based on areas without BCG coverage in the very large 1979 national survey sample.

Based on infection risk estimates from the five 5-yearly national surveys available from the Republic of Korea, the trend appears to be downward at 3.8 percent a year to a level in 1984 of 2 percent.

#### IV. DISCUSSION

Interpretation of this collection of risk and risk trend estimates must take into consideration the many potential sources of error, both in determining absolute levels of risk and in comparing risk estimates among countries or between periods within the same country. Some of these sources of error appear to have been adequately controlled, but the influence of other sources of error (identified and unidentified) must be acknowledged as unknown.

Tuberculin skin testing is a usefully accurate technique for separating groups at relatively high and low risk of future Mycobacterium tuberculosis disease, but is liable to varying degrees of error in identifying all the infected and specifying all the uninfected.

Sensitization to environmental mycobacteria is a source of nonspecific reactivity to tuberculin that is especially prevalent in tropical regions (Edwards and Edwards, 1960). There was evidence in reaction size distributions of a high prevalence of nonspecific sensitivity in several of the African and Asian populations surveyed, but distribution data for older groups or for bacteriologically confirmed cases were frequently available for locating the mode of the distribution of true infections and deriving their number by assuming the distribution to be symmetric.

There is underlying variability in tuberculin reactivity as evidenced by the variation between simultaneously applied identical tests in the same individual (Chaparas et al, 1985). Although this reduces precision of estimation, it is not expected to introduce systematic error.

A variety of antigen preparations and dosages were used, and these differed between countries and within countries over time. Estimating the number infected by using the symmetry of the distribution of true reactions about the mode would adjust for differences in specific biological potency between antigen preparations, but would not be effective in controlling for large differences in specificity of antigens.

The Mantoux method of administration was virtually always used, and all the selected studies measured induration of reaction. Techniques of measurement varied somewhat, but almost always consisted of transverse measurement at 48 or 72 hours after administration.

Waning of tuberculin reaction size over time, and complete reversion to zero reactivity are recognized to occur (Sutherland, 1971). To the extent reversions have occurred in a population, tuberculin test prevalence at a point in time will underestimate the proportion who were ever infected. But limiting to groups of about the same low average age may have minimized and kept comparable the effects of cumulative reversion.

Boosting is a potential problem in survey designs that retest the same individuals (Styblo, 1984). Boosting is the effect of an initial tuberculin test which, although it elicits slight or no reactivity itself, apparently acts as an antigenic stimulus able to recall waned or reverted preexisting mycobacterial sensitivity so that a subsequent test does elicit reactivity that is indistinguishable from conversion caused by M. tuberculosis transmission in the interval between the tests. Although some of the data compiled in this project came from surveys using a repeated testing design, the problem of boosting was avoided by selecting age groups too young to have been tested in earlier rounds.

BCG vaccination, a major component of tuberculosis control activities in developing countries, compromises the usefulness of tuberculin testing as a means to survey risk of infection. Tuberculin skin testing cannot distinguish between sensitivity caused by BCG vaccination and sensitivity caused by natural infection with M. tuberculosis, because BCG vaccination, unlike nonspecific sensitization, causes reactions that are distributed with nearly the same mode and shape as reactions caused by true infections with M. tuberculosis. For this reason the risk of infection in the whole population can only be inferred from risk measured in the unvaccinated.

In some of the countries for which survey data were selected, the policy was to vaccinate indiscriminantly without using tuberculin testing to select the as yet uninfected. In these countries, surveys limited to children without evidence of vaccination are subject to bias only to the extent that children selected for vaccination, either by choice or by local availability, are at a different risk of infection than the unvaccinated. This is a potentially severe source of bias if BCG coverage is very high.

In some of the surveys selected for this project, none of those surveyed had been vaccinated, either because BCG had not been introduced into that locale, or because BCG was withheld by policy until a later age. In other surveys, the unvaccinated were identified by physical examination to detect the characteristic scar. Levels of coverage by BCG were very low in some surveys, in which case representation of the whole population by the unvaccinated would be unaffected. In a few surveys, BCG coverage was so high that representation was demonstrably affected.

In other countries, the policy was to tuberculin test and vaccinate only those with weak or no reactions. In this situation, a later survey among children without BCG scars would be liable to severe overestimation bias. This was avoided by selecting age groups before the age of eligibility for selective vaccination programs in effect in a particular country.

Failure to test and measure every person in a survey sample can introduce a bias if those not reached by the survey are at a different risk of infection than those who were reached. A severe bias might also result from failure to measure a high proportion of those tested, because whether a survey participant returned for reading could well depend on whether there had been a reaction. Surveys with severe problems of this kind were not selected. Adequate coverage was a criterion for selection by this project, and was high in most of the surveys selected.

Although the average annual risk for a group lies in the calendar interval from birth to the time of the survey, it need not be at the interval midpoint. If the risk is changing substantially over time, as data from several of the surveys indicate, or varies with age, as has been suggested by the work of Sutherland (1976), and as might be expected in some of the diverse cultures surveyed, a complex model is necessary to estimate the risk at a particular time (Styblo, 1969). Limiting to young, narrow age groups was used as a way to bracket risk estimates within usefully narrow time intervals.

Average annual risk estimates for the same populations surveyed in different years are connected with straight lines in Figure 2 to indicate the average trend that would result in the observed risk estimates. An exponential trend is to be expected in the situation of a long-term constant ratio of the number of new cases produced by each existing case (Sutherland, 1976). However, without data for intermediate years, it is difficult to know the true shape of a trend line. In some instances where more than 2 surveys were available, the trend appeared to follow a fairly constant geometric decrease, but in other instances, the trend appeared to have changed over time.

Many survey reports presented more detailed analyses of infection risk and its trend, but for the purposes of this review, methods were adopted that could be simply and consistently applied to all data. In the few instances of large disagreement, the present analysis may be in error.

## V. CONCLUSIONS

This attempt to review and compile data available from major surveys carried out in developing countries since 1975 falls far short of providing a comprehensive picture of tuberculous infection risk and trend in the entire developing world. The assembled data cannot safely be generalized beyond the specific target populations surveyed. However, the assembled data do provide several recent objective examples of apparent progress and lack of progress in tipping the balance against the tuberculosis epidemic.

The review found that since 1975 many large scale surveys have been successfully carried out in the developing countries, with impressive attention to correct procedure and technique.

In most of the countries repeatedly surveyed in whole or in part, risk of infection does appear to have diminished. In some of these countries, the decline appears to have been substantial, and current levels of risk have become very low in a few countries.

In other countries, however, rates of infection appear to remain high and there is no indication of decrease.

It may be speculated that general improvements in living standards and tuberculosis control programs are having an effect against the tuberculosis epidemic in some parts of the developing world as it already has in the more developed countries. In the rest of the developing world, however, very little is known of the current magnitude and trend of the risk of tuberculous infection.

## REFERENCES

- Bleiker, M.A. & Styblo, K. (1978) The annual tuberculosis infection rate and its trend in developing countries. Bulletin of the International Union Against Tuberculosis, 53, 295-298.
- Comstock, G.W., Furculow, M.L., Greenburg, R.A., Grzybowski, S., Maclean, R.A., Baer, H. & Edwards, P.Q. (1971) The tuberculin skin test. American Review of Respiratory Disease, 104, 769-775.
- Chaparas, S.D., Vandiviere, H.M., Melvin, I., Koch, G. & Becker, C. (1985) Tuberculin test. Variability with the Mantoux procedure. American Review of Respiratory Disease, 132, 175-177.
- Edwards, P.Q. & Edwards, L.B. (1960) Story of the tuberculin test from an epidemiologic viewpoint. American Review of Respiratory Disease, 81, suppl., 1-47.
- Nyboe, J. (1960) The efficacy of the tuberculin test. Bulletin of the World Health Organization, 22, 5-37.
- Pio, A. Epidemiology of tuberculosis (1984) Minerva Medica, 75, 507-517.
- Styblo K., Meijer, J. & Sutherland, I. (1969) The transmission of tubercle bacilli. Its trend in a human population. Tuberculosis Surveillance Research Unit Report No. 1. Bulletin of the International Union Against Tuberculosis, 62, 1-104.
- Styblo K. (1984) Epidemiology of tuberculosis. Infektionskrankheiten und ihre Erreger, VEB Gustav Fischer Verlag, Jena.
- Styblo K. The relationship between the risk of tuberculous infection and the risk of developing infectious tuberculosis. Bulletin of the International Union Against Tuberculosis, 60, 117-119.
- Sutherland I. (1971) The effect of tuberculin reversion upon the estimate of the annual risk of infection. Bulletin of the International Union Against Tuberculosis, 45, 115-118.
- Sutherland I. (1976) Recent studies in the epidemiology of tuberculosis, based on the risk of being infected with tubercle bacilli. Advances in Tuberculosis Research, 19, 1-63.

TABLE 1. ANNUAL RISK OF TUBERCULOSIS INFECTION IN DEVELOPING COUNTRIES  
BASED ON TUBERCULIN SURVEYS<sup>a</sup>

- I. WHO African Region (except Algeria)
- II. WHO Region of the Americas
- III. WHO South-East Asia Region
- IV. WHO Eastern Mediterranean Region (and Algeria)
- V. WHO Western Pacific Region

<sup>a</sup> See our web site at URL: [www.who.int/bulletin](http://www.who.int/bulletin)

TABLE 1. ANNUAL RISK OF TUBERCULOUS INFECTION IN DEVELOPING COUNTRIES  
BASED ON TUBERCULIN SURVEYS

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year	Age (midpt.)	Tested N	Inferred N	Prevalence (%)	Risk/year (%)
<u>Botswana</u>										
1981-82ab	Whole country	Sample survey	83.3	RT23 2TU, Mode=13	1982.0	0-14 ( 6.5)	257	...	8.17	1.30
1956-57c	Whole country	Sample survey	0	RT19-20-21 5 TU, Mode=16	1957.0	0-14 ( 6.6)	1450	472	32.55	5.79
<u>Burundi</u>										
1982-84a	Military recruits	All recruits	71.0	RT23 2TU, 9mm+	1983.5	18-22 (20.5)	912	195	21.38	1.17
1964b	Northern section of Muramya Province	Sample survey	56.4	RT23 2TU, 9mm+	1964.6	15-19 (17.5)	202	76	37.62	2.66
<u>Cameroon - Cameroon</u>										
1984a	Yaounde	Sample survey of schools	58.4	PPD 71 10U, 9mm+	1984.5	7- 9 ( 8.5)	860	46	5.35	0.64
1964b	Urban Yaounde	Sample survey	...	RT23 2TU, 9mm+	1964.3	5- 9 ( 7.5)	326	31	9.51	1.32
<u>Ethiopia - Ethiopia</u>										
1983a	Addis Ababa	Sample of schools	49.4	RT23 2TU, 10mm+	1983.9	6-15 ( 8.6)	1251	133	10.63	1.30
1977b	Southwest area	Selected villages	...	... 2TU, 10mm+	1977.5	6-10 ( 8.5)	185	52	28.11	3.81
1953-55c	Addis Ababa	BCG school campaign	0	...	1954.7	7-14 (11 )	...	...	37	4.1
<u>Gambia</u>										
1976ab	Whole country	Sample survey	0.7	RT23 2TU, Mode=15	1976.4	6-16 ( 9.4)	2397	399	16.65	1.92
1958-59c	Bathurst in 2 strata	Sample survey:	...	RT23 1TU, Mode=17	1959.1	5-14 ( 9.4)	...	...	16.38	1.89
	Urban stratum	at 50 per 1000					176	61	34.66	4.43
	Rural stratum	at 7 per 1000					400	61	15.25	1.74

## I. WHO African Region (except Algeria) - Region Africaine de l'OMS (sans Algerie) (continued)

Year	Area Represented	Survey design	BCC (%)	Antigen and Infection Criterion	Survey Year Midpt. Age (midpt.)	Tested N	Infect ed N	Preva- lence (%)	Risk/ Year (%)
<b>Lesotho</b>									
1981-82 <sup>a</sup>	Whole country	Sample survey	70.4	RT23 2TU, Mode=14	1982.0 0-9 ( 5.2)	158	...	10.13	2.03
1956-57 <sup>b</sup>	Whole country	Sample survey	0	RT19-20-21 5TU, Mode=17,18	1957.0 0-9 ( 4.9)	1101	142	12.90	2.78
1962-65 <sup>c</sup>	Rural area around Maseru	Sample survey	0	RT23 1TU, Mode=19	1963.8 0-9 ( 4.9)	10216	1362	13.14	2.83
<b>Nigeria</b>									
1977 <sup>a</sup>	Kazaire Emirate of Kano State	School survey	32.1	RT23 2TU, Mode=15	1977.4 4-13*( 7* )	...	...	15.0	2.29
<b>United Republic of Tanzania - Republique-Unie de Tanzanie</b>									
1983 <sup>a</sup>	Arusha Region	Sample survey	52.4	RT23 2TU, Mode=17	1983.6 7-14 (10.0)	3125	307	9.82	1.03
1984	Coast (Pwani) Region	of schools	37.6		1984.8 ( 9.6)	681	74	10.87	1.19
1984	Dar-es-Salam Region	in progress, 90% completed	45.0		1984.8 ( 9.7)	1338	137	10.24	1.11
1986	Iringa Region		57		1986.4 (10.1)	1723	156	9.05	0.94
1986	Dodoma Region		49		1986.9 (10.7)	1743	358	20.54	2.13
1987	Kigoma Region		49.1		1987.7 (10.2)	1188	143	12.04	1.25
1985	Kilimanjaro Region		52.4		1985.4 (10.2)	2324	111	4.78	0.48
1984	Lindi Region		31.3		1984.6 (10.3)	2665	240	9.01	0.91
1986	Mbeya Region		64		1986.5 (10.9)	1621	193	11.91	1.16
1986	Morogoro Region		67		1986.6 (11.0)	865	158	18.27	1.82
1984	Mtwara Region		41.8		1984.5 ( 9.7)	629	122	19.40	2.20
1985	Mwanza Region		...		1985.8 (10.2)	2420	...	9.5	0.97
1986	Rukwa Region		67		1986.4 (10.8)	385	47	12.21	1.20
1984	Ruvuma Region		46.1		1984.5 (10.4)	1412	140	9.92	1.00
1985	Shinyanga Region		...		1985.4 ( 9.8)	3666	...	11.6	1.25
1985	Singida Region		...		1985.7 (10.5)	985	...	16.3	1.68
1987	Tabora Region		...		1987.7 (11.2)	1794	...	8.3	0.77
1983	Tanga Region		50.1		1983.8 (10.4)	2462	238	9.67	0.97
1983-87	(18 Area weighted average)				1985.6 (10.3)	30982	...	10.9	1.11
1979 <sup>b</sup>	Shinyanga Region, except Shinyanga Distr.	Sample survey	50.9	RT23 2TU, 10m+	1979.5 6-14 (10.0)	1817	356	19.59	2.16
1978 <sup>d</sup>	Dodoma Region	Sample survey	63.3	RT23 2TU, Mode=17	1978.9 6-14 (10.2)	1329	166	12.49	1.30
1977 <sup>e</sup>	Tanga Town and nearby rural area	Sample survey of schools	79.0	RT23 2TU, Mode=17	1977.1 7-14 ( 9.7)	383	33	8.62	0.92

## II. WHO Region of the Americas - Region des Ameriques de l'OMS

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt. Age (midpt.)	Tested		Infection		Prevalence (%)	Risk/Year (%)
						N	N	N	N		
<b>Argentina - Argentine</b>											
1974-78ab	Whole country	Sample survey	...	RT23 2TU, 10mm+	1976.5	6-7 (6.5)	26902	970	3.61	0.56	
1983c	Areas around	Sample survey	81.6	.... 10mm+	1983.5	5-8 (7.0)	325	...	1.8	0.26	
1979-80c	Santa Fe City		63.7	.... 10mm+	1980.0	6-7 (7.0)	443	...	2.0	0.29	
1967-68c			31.6	.... 10mm+	1968.0	5-8 (7.0)	1221	...	8.8	1.31	
1979-80c	Santa Fe City	All schools	57.7	.... 10mm+	1980.0	6-7 (7.0)	2125	...	1.6	0.23	
1974-75c			12.9	.... 10mm+	1975.0	6-7 (7.0)	3590	...	1.6	0.23	
1967-68c			...	.... 10mm+	1968.0	6-7 (7.0)	3196	...	3.9	0.57	
1960-61d	Resistencia Province	Sample survey	Rate	RT25 1TU, 10mm+	1961.0	5-9 (7.5)	1259	49	3.89	0.53	
<b>Brazil - Brasil</b>											
1983a	Rio Grande Do Sul Province	Sample survey of school enterers	0	RT23 2TU, 10mm+	1983.5	7-8 (7.3)	11880	333	2.80	0.39	
By Provincial zone:											
Metropolitan											
Southern rural											
Northern rural											
1980b	Capitals of States and Territories by Region:	All school enterers	...	...	1980.5	... (7.7)	...	...	...	...	
North											
Northeast											
Southeast											
Sao Paulo State											
South											
East Central											
1970-73b	Capitals of States and Territories by Region:	All school enterers	...	RT23 2TU, 10mm+	1972.0	... (7.7)	...	...	...	...	
North											
Northeast											
Southeast											
Sao Paulo State											
South											
East Central											

## III. WHO South-East Asia Region - Region de l'Asie du Sud-Est de l'OHS

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt.	Age (midpt.)	Tested		Infec- Preva- ted lence		Risk/ Year (%)	
							N	N	(%)	(%)		
<b>India - Inde</b>												
1979 <sup>a</sup> 1974-75de	Dodballapur in Bangalore District	Sample survey repeated in same villages	14.1 36	RT23 1TU, 12mm+ RT23 1TU, 10mm+	1979.5 1974.8	0-4 ( 2.5)	5203 3805	128 98	2.46 2.58	0.99 1.04		
1977-78 <sup>b</sup> 1961-63 <sup>a</sup>	3 Subdistricts in Bangalore District	Repeated sample surveys	... 1.5	RT23 1TU, 10mm+	1977.9 1962.2	0-4 ( 2.5)	1492 7981	36 166	2.41 2.08	0.97 0.84		
1972-73 <sup>f</sup> 1960-61 <sup>cfi</sup>	Tumkur District in South India	Sample survey repeated in same 62 villages	... ...	RT23 1TU, 10mm+	1973.0 1960.9	0-4 ( 2.5)	4277 3788	115 155	2.69 4.09	1.08 1.66		
1978 <sup>g</sup>	Kashmir Valley	Sample survey	0.2	PPD-S 3TU, 12mm+	1978.7	0-4 ( 2.5)	2448	56	2.29	0.92		
1972 <sup>h</sup>	Villages in: Kashmir Kulu Valley Lohaghat Pithoragarh Agra Haryana Rajasthan	Selected for altitude contrast	<5	PPD-S 5TU, 12mm+	1972.8	1-4 ( 3.0)	679 503 381 290 530 612 76	... ... ... ... ... ... ...	4.0 5.2 3.1 4.1 5.2 4.4 1.3	1.4 1.8 1.0 1.4 1.8 1.5 0.4		
1968-71 <sup>h</sup>	Northern Chingelput	Selected for BCG trial	<5	PPD-S 5TU, 12mm+	1969.9	1-4 ( 3.0)	27520	...	5.1	1.7		
<b>Indonesia - Indonesie</b>												
1983 <sup>aefgh</sup> 1978 1972	Tangerang Regency in West Java	Sample survey repeated in same schools	18.5 22.1 1.9	RT23 2TU, 10mm+	1983.6 1978.4 1972.2	7-10 ( 8.8) 7-10 ( 8.8) 7-10 ( 9.3)	1549 1649 1371	286 440 497	18.46 26.68 36.25	2.29 3.47 4.73		
1986 <sup>hh</sup> 1981 1976	Sambas in West Kalimantan	Selected as BCG-free area	0	RT23 2TU, 10mm+	1986.8 1981.8 1976.9	7-10 ( 8.8) 7-10 ( 8.6) 7-10 ( 8.6)	3839 2181 1655	576 310 391	15.00 14.21 23.63	1.83 1.77 3.09		
1986 <sup>hh</sup> 1981 1976	Padang Pariaman in West Sumatra	Selected as BCG-free area	0	RT23 2TU, 10mm+	1986.9 1981.7 1976.5	7-10 ( 8.7) 7-10 ( 8.9) 7-10 ( 8.6)	1986 2501 1124	472 363 193	23.77 14.51 17.17	3.07 1.75 2.17		
1984 <sup>gh</sup> 1979 <sup>cgh</sup>	Malang in East Java	Selected as BCG-free area	0	RT23 2TU, 10mm+	1984.5 1979.5	7-10 ( 8.3) 7-10 ( 8.9)	1406 1122	75 86	5.33 7.66	0.66 0.89		
1985 <sup>cgh</sup> 1980 1975	Ogan Kaming Ilir in South Sumatra	Selected as BCG-free area	0	RT23 2TU, 10mm+	1985.8 1980.2 1975.6	7-10 ( 8.9) 7-10 ( 8.5) 7-10 ( 8.9)	4840 4839 2425	1309 1358 701	27.05 28.06 28.91	3.48 3.80 3.76		

111. WHO South-East Asia Region - Region de l'Asie du Sud-Est de l'OHS (continued)

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt.	Age (midpt.)	Tested N	Infect ed N	Preva- lence (%)	Risk- year (%)
1985 <sup>cgh</sup>	Gowa in South Sulawesi	Selected as BCG-free area	0	RT23 2TU, 10mm+	1985.5	7-10 ( 8.9)	4001	1198	29.94	3.92
1980					1980.5	7-10 ( 8.7)	3573	1076	30.11	4.03
1975					1975.8	7-10 ( 8.2)	1429	429	30.02	4.26
1983 <sup>cgh</sup>	Stabat Langkat in North Sumatra	Selected as BCG-free area	0	RT23 2TU, 10mm+	1983.2	7-10 ( 8.5)	1894	196	10.35	1.28
1978					1978.4	7-10 ( 8.9)	1125	165	14.67	1.77
1982 <sup>cgh</sup>	Hulu Sengai Tengah in South Kalimantan	Selected as BCG-free area	0	RT23 2TU, 10mm+	1982.8	7-10 ( 9.0)	1577	424	26.89	3.42
1977					1977.3	7-10 ( 8.7)	1199	352	29.36	3.92
1984 <sup>gh</sup>	Pati in Central Java	Selected as BCG-free area	0	RT23 2TU, 10mm+	1984.5	7-10 ( 8.5)	2938	403	13.72	1.72
1979					1979.5	7-10 ( 8.9)	2197	381	17.34	2.12
1974					1974.5	7-10 ( 8.1)	1070	143	13.36	1.76
1983 <sup>B</sup>	Langsa in D.I. Aceh	Selected as BCG-free area	0	RT23 2TU, 10mm+	1983.5	7-10 ( 9 )	...	...	24.4	3.06
1964-65 <sup>dh</sup>	Rural area in East Java	Sample survey	...	RT23 2TU, 10mm+	1965.0	5- 9 ( 7.5)	1633	191	11.70	1.64
<u>Thailand - Thaïlande</u>										
1977-79 <sup>a</sup>	Whole country	Sample survey	39.1	...	1978.5	0-14 ( 7.5)	...	...	15.2	2.17
1960-64 <sup>bd</sup>	Bangkok and Chiangmai Province	Sample survey	...	...	1962.5	10-14 (12.5)	...	...	29	2.7
1954 <sup>C</sup>	6 localities in Chiangmai and Kanchanaburi Provinces	Selected unvaccinated areas	0	... 5TU, 10mm+	1954.9	8-12 (10.5)	1578	371	23.51	2.52

## IV. WHO Eastern Mediterranean Region (and Algeria) - Region de la Mediterranee Orientale de l'OMS (et Algerie)

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt.	Age (midpt.)	Tested		Prevalence (%)	Risk/year (%)
							N	N		
<b>Algeria - Algeria</b>										
Whole country in 6 economic zones										
1985d	Zone 2 Tizi Ouzou and Tlemcen	Second survey (commencing)	67.2	RT23 2TU, 10mm+	1985.5	8 ( 8.5)	2378	55	2.31	0.27
1984d	Zone 5 & 6	First survey	59.1		1984.5		2125	64	3.01	0.36
1983ab	Zone 4 incl. Batna	(nearly complete)	68.6		1983.9		1144	51	4.46	0.54
1982ab	Zone 3 incl. Guelma		66.3		1982.9		1067	45	4.22	0.51
1981ab	Zone 1 incl. Mascara		63.3		1981.9		1334	71	5.32	0.64
1980ab	Zone 2 Tizi Ouzou and Tlemcen		44.6		1980.9		1844	71	3.85	0.46
1980-84	(5 Area weighted average)				1982.8		7514	302	4.02	0.48
1981ab	Blida	Survey repeated in some schools	71.6	IP48 10TU, 6mm+	1981.5	8 ( 8.5)	1117	...	6.18	0.75
1976ab	Blida				1976.5	6-10 ( 8.5)	262	...	8.4	1.03
1949-52C	Whole country	Mass BCG Campaign	...	IP48 5TU, 6mm+ IP48 10TU, 6mm+ Moro Patch, 3+ papules	1951.1	8 ( 8.5)	110547	...	31.2	4.30
<b>Afghanistan</b>										
1982ac	Whole country	Sample survey	...	RT23 1TU, 8mm+	1982.5	5- 9 ( 7.5)	881	208	23.61	3.53
1963bc	Kabul	Selected schools	...	RT23 1TU, 8mm+	1963.5	7-12 (10.0)	19006	4322	22.74	2.55
	Jalalabad					7-12 (10.0)	3492	1083	31.01	3.64
	Mazar-e-Sharif					7-12 (10.0)	3170	1019	32.15	3.80
	Kandahar					7-12 (10.0)	4079	1586	38.88	4.80
	Pulkitumry					13-15 (14.5)	1191	384	32.24	2.65
	(5 Area weighted average)					7-15 (10.2)	30938	8394	27.13	3.06
<b>Bahrain - Bahrain</b>										
1981a	Whole country	All schools	0	RT23 1TU, 6mm+	1981.4	6- 7 ( 7.0)	6151	86	1.40	0.20
1969a	Whole country	Sample survey	...	RT23 1TU, 6mm+	1969.5	6- 7 ( 7.0)	897	55	6.13	0.90

## IV. WHO Eastern Mediterranean Region (and Algeria) - Region de la Méditerranée Orientale de l'OHS (et Algérie) (continued)

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt. Age (midpt.)	Tested N	Infect ed N	Preva- lence (%)	Risk/ Year (%)
<u>Kuwait - Kuwait</u>									
1981 <sup>a</sup>	Whole country	School campaign	0	RT23 ...	1981.5	22710	261	1.15	0.22
1980					1980.5	16149	101	0.63	0.12
1979					1979.5	22674	283	1.25	0.24
1978					1978.5	20843	338	1.62	0.31
1977					1977.5	17444	266	1.52	0.29
1976					1976.5	9018	201	2.23	0.44
1975					1975.5	7665	100	1.30	0.29
1974					1974.5	6722	55	0.82	0.20
1973					1973.5	6363	80	1.26	0.29
1972					1972.5	2258	38	1.68	0.36
1972-81	(10 Year weighted average)				1978.4	131846	1723	1.31	0.26
1962-63 <sup>a</sup>	Whole country	Sample survey	...	...	1963.0	...	...	10.74	1.50
<u>Libya Arab Jamahiriyah - Jamahiriyah Arabe Libyenne</u>									
1976-77 <sup>a</sup>	Whole country East Libya	Sample survey	63.5	RT23 2TU, Mode=16,17	1977.0	1827	35	1.92	0.26
1959 <sup>ab</sup>	East Province Cyrenaica	Sample survey	...	RT23 1TU, Mode=18,19	1959.7	361	60	16.62	2.39
1954 <sup>ac</sup>	Ben Walid	Selected area	0	RT19-20-21 5TU, Mode=18,19	1954.5	188	...	24.5	3.68
<u>Pakistan</u>									
1974-78 <sup>abd</sup>	Whole country except Baluchistan	Sample survey	7.8	RT23 1TU, 10mm+	1976.5	2289	298	13.02	1.84
1961-62 <sup>a-d</sup>	Karachi, Rawalpindi, and Lahore	Sample survey	42.7*	RT23 1TU, 10mm+	1962.0	769	178	23.15	3.45
1949-54 <sup>be</sup>	Provincial capitals Other urban areas Rural areas (3 Area median)	Mass BCG campaign	0	RT22 5TU, 5mm+	1954.1	...	...	47.5	5.69
								44.5	5.21
								41.0	4.68
								44.5	5.21

## IV. WHO Eastern Mediterranean Region - Region de la Méditerranée Orientale de l'OMS (continued)

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt.	Age (midpt.)	Tested		Prevalence		Risk/	
							N	N	(%)	(%)	Year	(%)
<u>Syrian Arab Republic - République Arabe Syrienne</u>												
1983 <sup>a</sup>	Aleppo	Survey repeated in some 1978 survey schools	14.1	RT23 2TU, 10mm+	1983.8	6 ( 6.5)	1586	31	1.95	0.30		
1978 <sup>b</sup>	Aleppo	Sample survey of schools	8.5	RT23 2TU, 10mm+	1978.8	6 ( 6.5)	1845	31	1.68	0.26		
1950 <sup>ad</sup>	Aleppo	BCG school campaign	0	Moro patch, 3+ papules	1950.3	6 ( 6.5)	...	...	15.3	2.52		
1983 <sup>e</sup>	Homs	Sample survey of schools	17.2	RT23 2TU, 10mm+	1983.8	6 ( 6.5)	1182	11	0.93	0.14		
1950 <sup>d</sup>	Homs	Mass BCG campaign	0	Moro patch, 3+ papules	1950.6	6 ( 6.5)	...	...	10.4	1.68		
1960 <sup>c</sup>	Districts of Homs and Damascus, excluding City of Damascus	Sample survey	9*	RT23 1TU, 10mm+	1960.8	5- 9 ( 7.5)	387	21	5.43	0.74		

## V. WHO Western Pacific Region - Region du Pacifique Occidental de l'OMS

Year	Area Represented	Survey design	BCG (%)	Antigen and Infection Criterion	Survey Year Midpt. Age (midpt.)	Tested N	Infectious N	Prevalence (%)	Risk/Year (%)
<u>China - Chine</u>									
1979ab	Areas without BCG, in whole country except Taiwan	266 of the 888 clusters in the national sample	0	RT23 2TU, 6mm+	1979.5 7 ( 7.5)	10000*	...	7.3	1.01
<u>Malaysia - Malaisie</u>									
1976-77a	Whole country	Sample survey	81-86	RT23 1TU, Mode=15	1977.0 1- 9 ( 5.5)	1429	29	2.03	0.37
<u>Republic of Korea - Republique de Coree</u>									
1985ab	Whole country	Sample survey	65.9	RT23 1TU, 10mm+	1985.5 0- 4 ( 2.8)	1420	77	5.42	1.97
1980cd	Whole country	Sample survey	50.0	RT23 1TU, 10mm+	1980.5 3m- 4 ( 2.1)	1310	...	4.89	2.36
1975c	Whole country	Sample survey	48.2	RT23 1TU, 10mm+	1975.5 3m- 4 ( 2.1)	1871	...	4.8	2.32
1970c	Whole country	Sample survey	...	RT23 1TU, 10mm+	1970.5 3m- 4 ( 2.1)	...	...	8.5	4.14
1965cef	Whole country, 2 strata Urban stratum Rural stratum	Sample survey at 1 per 1040 at 1 per 2220	6.9	RT23 1TU, 10mm+	1965.5 0- 4 ( 2.5)	..	..	10.25	4.23
<u>Philippines</u>									
1981-83ab	Whole country	Sample survey	38.9	RT23 1TU, Mode=14,15	1982.5 2m- 4 ( 2.1)	2038	78	3.83	1.84
<u>Samoa</u>									
1975a	Main island (Upolu)	Sample survey	72.6	RT23 1TU, 10mm+	1975.6 3m- 7 ( 2.8)	1824	25	1.37	0.49
1966-68bc	Whole country	BCG Campaign	...	RT23 1TU, 10mm+	1967.3 0- 9 ( 5.0)	1710	46	2.69	0.54
1961-63b	Whole country	All persons	...	RT23 1TU, 10mm+	1962.5 0- 9 ( 5.0)	1284	22	1.71	0.35

Symbols: \* Estimated .. Not applicable ... Not available