

WHO maternal death and near-miss classifications

Robert Pattinson,^a Lale Say,^b João Paulo Souza,^b Nynke van den Broek^c & Cleone Rooney^d on behalf of the WHO Working Group on Maternal Mortality and Morbidity Classifications

Reducing maternal mortality is the Millennium Development Goal 5. To reach this goal, countries need an accurate picture of the causes and levels of maternal deaths. Recent systematic reviews have shown that there are many inconsistencies in the way maternal deaths are classified and there is a lack of standard definitions and criteria for identifying severe maternal morbidity and near-miss cases.^{1,2}

WHO established a technical working group of obstetricians, midwives, epidemiologists and public health professionals from developing and developed countries to develop a maternal death classification system.

The group established three principles for its work. First, the classification must be practical and understood by its users (clinicians, epidemiologists and programme managers). Second, underlying causes must be exclusive of all other conditions; as in the *International Statistical Classification of Diseases and Related Health Problems* (ICD), the underlying cause is the disease or injury which initiated the sequence of events leading directly to death, or the circumstances of the accident or violence which produced the fatal injury. Third, the new classification system should be compatible with and contribute to the 11th revision of the ICD. Incorporating this maternal death classification into the ICD will encourage consistent use in both death certificates and confidential enquiries into maternal deaths, and improve the comparability of data.

The proposed maternal death classification system was sent to more than 40 individual reviewers and the International Federation of Gynecology and Obstetrics, the United Nations Children's Fund (UNICEF), the United Nations Population Fund (UNFPA)

and national professional organizations including the Royal College of Obstetricians and Gynaecologists, the American College of Obstetricians and Gynecologists and the Canadian College of Obstetricians and Gynaecologists. Revised after this feedback, the second version was tested on eight databases of maternal deaths: national databases from Colombia, Jamaica and South Africa, other databases from Kenya, Malawi and Zimbabwe and verbal autopsy data from Afghanistan and Nigeria. All sites found the classification workable and useful.

The new WHO classification of cause of maternal death has a simple structure to facilitate tabulation: group, disease category and individual underlying causes. The group includes three categories: direct maternal deaths, indirect maternal deaths and "unanticipated complications of management". This addition makes it possible to track trends in iatrogenic disease as, for example, related to caesarean sections. Underlying causes are clearly separated from conditions contributing to fatal outcomes. Finally, the working group decided to classify suicide in pregnancy, deaths from puerperal psychosis and postpartum depression in the category of direct maternal deaths.³

The working group also reached consensus on how to define maternal near miss – "a woman who nearly died but survived a complication that occurred during pregnancy, childbirth or within 42 days of termination of pregnancy".⁴ Signs of organ dysfunction that follow life-threatening conditions are used to identify maternal near misses so that the same classification of underlying causes is used for both maternal deaths and near misses. This consistency and a set of near-miss

indicators enables assessments of the quality of care provided to pregnant women.

The WHO technical working group recommends that the new maternal death classification system be adopted by all countries and the maternal near-miss approach be considered in national plans for improving maternal health. By using the same classifications, reliable comparisons can be made within and between countries and regions. Applying this classification should help to identify the health system shortfalls that countries need to address in order to reduce complications and fatal outcomes of pregnancy and childbirth. ■

Acknowledgements

Available at: <http://www.who.int/bulletin/volumes/87/10/09-071001/en/index.html>

References

1. Khan KS, Wojdyla D, Say L, Gulmezoglu AM, Van Look PFA. WHO analysis of causes of maternal deaths: a systematic review. *Lancet* 2006;367:1066-74. PMID:16581405 doi:10.1016/S0140-6736(06)68397-9
2. Say L, Pattinson RC, Gülmezoglu AM. WHO systematic review of maternal morbidity and mortality: the prevalence of severe acute maternal morbidity (near miss). *Reprod Health* 2004;1:3. PMID:15357863 doi:10.1186/1742-4755-1-3
3. *Report on the World Health Organization Working Group on the Classification of Maternal Deaths and Severe Maternal Morbidities*. Geneva: World Health Organization; 2009.
4. Say L, Souza JP, Pattinson RC. Maternal near miss – towards a standard tool for monitoring quality of maternal health care. *Best Pract Res Clin Obstet Gynaecol* 2009;23:287-96. PMID:19303368 doi:10.1016/j.bpobgyn.2009.01.007

^a MRC Maternal and Infant Health Care Strategies Research Unit, University of Pretoria, Gauteng, South Africa.

^b Department of Reproductive Health and Research, World Health Organization, 20 avenue Appia, 1211 Geneva 27, Switzerland.

^c Liverpool School of Tropical Medicine, Liverpool, England.

^d Office for National Statistics, London, England.

Correspondence to João Paulo Souza (e-mail: souzaj@who.int).

Acknowledgements

The following individuals were among the members of the expert group that developed the identification criteria for maternal near miss: Affette McCaw-Binns, Anoma Jayathilaka, Buyanjargal Yadamsuren, Cleone Rooney, João Paulo Souza, José Guilherme Cecatti, Lale Say, Linda Bartlett, Mary Ellen Stanton, Mohamed Cherine Ramadan, Nynke van den Broek, Robert C Pattinson, Rogelio Gonzalez, Veronique Filippi. We thank Ahmet Metin Gülmezoglu, Jelka Zupan, and Vicky Camacho for their contribution. This work was funded by USAID and UNDP/UNFPA/WHO/World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP).