

# Screening eye diseases in babies: an Italian experience on 5000 healthy, consecutive newborns

Roberto Perilli<sup>1</sup>, Modesto Lanci<sup>2</sup>, Antonino Romano<sup>3</sup>, Laura Sabatini<sup>4</sup> and Paola Fusilli<sup>4</sup>

<sup>1</sup>UOSD Oftalmologia Sociale Territoriale, AUSL Pescara, Pescara, Italy

<sup>2</sup>UOC Oculistica, Ospedale "Spirito Santo", Pescara, Italy

<sup>3</sup>UOC Oculistica, Ospedale Pediatrico Bambino Gesù, Rome, Italy

<sup>4</sup>UOC Neonatologia, Ospedale "Spirito Santo", Pescara, Italy

## Abstract

**Introduction.** Visual performance of eyes with congenital pathologies is conditioned by an early diagnosis. Families having problems in accessing health services risk to delay or miss both an early diagnosis and an early treatment and amblyopia (lazy eye) prevention.

**Methods.** In our hospital, all full-term, healthy newborns are thoroughly examined by an ophthalmologist in the maternal ward, 1 to 3 days after birth.

**Results.** Among the first 5000 newborns examined, a high incidence of congenital pathologies compared to international literature was reported, with differences between Caucasians and non-Caucasians.

**Conclusion.** Performing an early in-hospital thorough eye examination in all newborns as a screening would be an effective way to miss none and to start an early and effective pathway of disease treatment.

## Key words

- congenital eye diseases
- congenital cataract
- congenital glaucoma
- red reflex test
- newborn eye examination

## INTRODUCTION

The rapid postnatal increase of visual acuity [1, 2] requires an early identification of congenital eye pathologies. The most diffuse and sight-threatening pathologies (cataract and glaucoma) are often diagnosed during a pediatric examination later in the infant's life, or by an altered red reflex in photographs, thus delaying interventions for disease treatment and amblyopia prevention. Chan *et al.* [3] reported that visual outcomes are better in children undergoing surgery before 6 weeks of age in unilateral cataracts and before 10 weeks of age in bilateral ones. Mandal *et al.* [4] reported that the majority of congenital glaucomas (about 60%) are diagnosed by the age of 6 months, and 80% are diagnosed within the first year of life.

In 2008, the American Academy of Pediatrics (AAP) [5] provided the pediatricians with guidelines on the Red Reflex Test (RRT): this must be performed, possibly at birth, by a trained pediatrician in a darkened room, projecting the light of a direct ophthalmoscope into both eyes from approximately 18 inches away, and checking that a red reflex comes back from both eyes.

In 2012, a non-institutional survey [6] reported that in Italy in only 64% of the maternal wards, and in none with more than 2500 births per year, the RRT was performed by a pediatrician in all newborns at birth.

In this paper, we demonstrate how an ophthalmologist can easily integrate the RRT with an examination of both the anterior segment and the fundus of the newborns' eyes in a handful of seconds, thus providing a prompt diagnosis and allowing the ill babies to start the therapeutic pathway as soon as possible.

## MATERIALS AND METHODS

Since June 2011, at the maternity ward of the "Spirito Santo" Hospital in Pescara, Italy, the eye examination is part of an institutional and parent-approved series of routine clinical ones in all full-term, healthy newborns, 1 to 3 days after birth.

In February 2014, 5000 consecutive newborns (Caucasian: 2427 males and 2334 females; non-Caucasian: 116 males and 123 females; see *Table 1* for distribution by sex and ethnicity) had received the test by an ophthalmologist. Pupils are dilated with 0.5% tropicamide drops (Visumidriatic 0.5%, Visupharma, Italy), gently pressing the inner lid canthus in order to reduce systemic absorption. Then, in a darkened room, the ophthalmologist, with a direct ophthalmoscope (Welch-Allyn, USA), checks the red reflex of both eyes from an arm's distance (according to the AAP guidelines), and thereafter adds a closer exam of the anterior segment (focusing the iris plane) and posterior pole (focusing

**Table 1**  
Sample distribution by ethnicity and sex

Ethnicity	Males	Females	Total (%)
Total sample	2547	2457	5000 (100)
Caucasian	2427	2334	4761 (95.2)
Non Caucasian	116	123	239 (4.8)
African	79	85	164(3.3)
Chinese	23	17	40 (0.8)
Hispanic	14	19	33 (0.7)
Indian	0	2	2 (0.04)

the retina), at a few centimeters' distance. Moreover, he disposes of a ruler, to measure the corneal diameters in the suspect of buphthalmos in congenital glaucoma.

Tests are routinely performed with the parents' informed consent and data treated in compliance with the Declaration of Helsinki. The present note is a report of routine clinical practice. All pathologic infants are sent to the Bambino Gesù Children Hospital in Rome, for therapy and for a thorough study of both the newborns' clinical condition and the parents' health status.

## RESULTS

No newborn suffered from adverse effects to tropicamide. Pathologies are summarized in Table 2. Cataracts (n = 7) were both bilateral and involving the visual axis (n = 4), or unilateral and sparing the visual axis (n = 3). Glaucomas (n = 3) were all bilateral.

In addition, one case of Primary Hyperplastic Vitreous Persistence (PHVP), one case of angle dysgenesis and one case of inferior coloboma of the iris, retina and choroid were reported.

Embryonic vitreous remnants were frequent: pupillary strands (n = 679) and/or (multi)focal ectropion uveae (n = 128, mainly in the superior pupillary margin). Cases with ectropion uveae were given particu-

lar attention to other signs of local (glaucoma) and/or systemic (neurofibromatosis or other neural crest cell disorders) disease, always absent, and reported to the newborn's pediatrician.

Retinal haemorrhages were found in 79 newborns, all of which born by vaginal delivery.

## DISCUSSION

Tropicamide 0.5% confirmed being a safe drug [7]. International literature reports a wide range of incidence as for congenital pathologies: congenital cataract: 0.1-1.5/1000 in general [8], 0.3/1000 (2 out of 3 bilateral) in the United Kingdom [9]; congenital glaucoma: 0.1/1000 in developed countries, 0.8/1000 in Slovakian Roms, 0.4/1000 in the Middle East, 0.31/1000 in Andhra Pradesh, India [10]; coloboma of the iris and retina: 0.48/1000 in Minnesota, USA, 0.037/1000 in Hungary and 0.08/1000 in Scotland [11]. Retinal haemorrhages involving the macula can interfere with the natural development of visual function in the first days or weeks [12].

In our series, the incidence of pathologies ranks high in worldwide reports, possibly because all newborns were screened, thus avoiding missing cases.

With limitations owing to the small sample, there was a significant difference between Caucasian and non-Caucasian newborns: cataract's incidence was 1.05/1000 in Caucasians and 8.37/1000 in non-Caucasians ( $p < 0.005$ ); glaucoma's incidence was 0.21/1000 in Caucasians and 8.37/1000 in non-Caucasians ( $p < 0.001$ ).

## CONCLUSIONS

We believe that our results raise important problems of health policy: missing the opportunity of a thorough eye exam or – at least – a RRT at birth, particularly in babies who may have a delayed and/or difficult approach to (public) assistance, increases the risk of not having their pathologies identified and treated in due time. This appears to be important particularly for non-Caucasian babies, showing a higher incidence of pathologies.

**Table 2**  
Sample distribution by ethnicity, sex and pathology

Type of pathology	n	Male incidence (%)	n	Female incidence (%)	n	Total incidence (%)
Cataract in Caucasians	2	0.82	3	1.29	5	1.05
Cataract in non Caucasians	0	0	2	16.26	2	8.37
Glaucoma in Caucasians	0	0	1	0.43	1	0.21
Glaucoma in non Caucasians	0	0	2	16.26	2	8.37
PHVP ^*	0	0	1	0.41	1	0.20
Angle dysgenesis*	0	0	1	0.41	1	0.20
Coloboma*	0	0	1	0.41	1	0.20
Pupillary strands*	367	144.32	312	126.98	679	135.80
Ectropion uveae*	96	37.75	32	13.02	128	25.60
Retinal haemorrhages*	42	16.52	37	15.06	79	15.80

^PHVP: primary hyperplastic vitreous persistence;

\*Incidence % is calculated on the whole sample of 5000 newborns (2543 males and 2457 females) independently from ethnicity.

Aside of genetic factors, it is well known [13, 14] that a series of social ones is related to health (mainly infectious diseases) and mortality: income, education, occupation, marital status and so on. Moreover, in different ethnicities the prevalence of both systemic and ocular diseases (infectious and non-infectious) is heterogeneous, and the immunization status can be suboptimal [15]: this could account for subclinical or latent conditions predisposing to a pathologic development of eye structures.

Finally, from a strictly scientific viewpoint, it is very interesting to notice how a careful and close examination of structures like the iris and pupil (missed by the RRT) can show aspects of late intrauterine development of the eye: such structures are involved in the regression of the tunica vasculosa lentis (a membrane covering the lens and extending across the pupil, regressing before birth) [16]: the high incidence of pupillary strands and (multi)focal ectropion uveae, disappearing in the weeks/months after birth and never related to eye or systemic

pathologies, shows how late intrauterine processes can continue even after birth (what has been pathologically shown by the case of PHVP).

In our experience, the RRT can be easily integrated with a thorough eye examination and performed at the maternity ward by an ophthalmologist in a very short time, thus possibly widening the spectrum of clinical conditions being detected. In Italy, the eye examination at birth is not always provided as a routine procedure in hospital settings, but we believe it should be encouraged, since it fulfills both the 1968 Wilson and Jungner's [17] and the 2008 World Health Organization's [18] criteria for screenings.

#### Conflict of interest statement

None of the authors has any kind of conflict of interest.

Received on 1 April 2015.

Accepted on 17 September 2015.

## REFERENCES

- Chandna A. Natural history of the development of visual acuity in infants. *Eye* 1991;5:20-6.
- Lewis TL, Maurer D. Multiple sensitive periods in human visual development: evidence from visually deprived children. *Dev Psychobiol* 2005;46:163-83.
- Chan W, Biswas S, Ashworth JL, Lloyd IC. Congenital and infantile cataract: aetiology and management. *Eur J Pediatr* 2012;171(4):625-30.
- Mandal AK, Chakrabarti D. Update on congenital glaucoma. *Indian J Ophthalmol* 2011;59(Suppl. 1):S148-S157.
- American Academy of Pediatrics. Red Reflex Examination in Neonates, Infants and Children. *Pediatrics* 2008;122:1401-4.
- Cittadinanza attiva onlus. *Presentazione report "Percorso nascita". Percorso nascita: i centri più grandi offrono maggiori tutele. Necessario uniformare i servizi delle Regioni per garantire uguale accesso a prestazioni e screening neonatali.* 27 settembre 2012. Available from: [www.cittadinanzattiva.it/comunicati/salute/maternita-fertilita/3977-presentazione-report-percorso-nascita.html](http://www.cittadinanzattiva.it/comunicati/salute/maternita-fertilita/3977-presentazione-report-percorso-nascita.html).
- Ogut MS, Bozkurt N, Ozek E, Birgen H, Kazokoglu H, Ogut M: Effects and side effects of mydriatic eye drops in neonates. *Eur J Ophthalmol* 1996;6:192-6.
- Yi J, Yun J, Li Z-K, Xu C-T, Pan B-R. Epidemiology and molecular genetics of congenital cataracts. *Int J Ophthalmol* 2011;4(4):422-32.
- Rabi JS. Measuring and interpreting the incidence of ocular anomalies: lessons from a national study of congenital cataracts in the UK. *IOVS* 2001;4:1444-8.
- Mandal AK, Chakrabarti D. Update on congenital glaucoma. *Indian J Ophthalmol* 2011;59(Suppl. 1):S148-S157.
- Nakamura KM *et al.* Incidence, Ocular Findings and Systemic Associations of Ocular Coloboma: A Population-Based Study. *Arch Ophthalmol* 2011;129(1):69-74.
- Li LH, Li N, Zhao JY, Fei P, Zhang G, Mao G, Rychwalski PJ. Findings of perinatal ocular examination performed on 3573 healthy full term newborns. *Br J Ophthalmol*. 2013;97:588-91.
- Richardus JH, Kunst AE. Black-white differences in infectious disease mortality in the United States. *Am J Public Health*. 2001;91(8):1251-3.
- Sorlie PD, Backlund E, Keller JB. US Mortality by Economic, Demographic, and Social Characteristics: the National Longitudinal Mortality Study. *Am J Public Health* 1995;85(7):949-56.
- Canadian Collaboration for Immigrant and Refugee Health. Evidence-based clinical guidelines for immigrants and refugees. *CMAJ* 2011;183(12):E824-E925
- Pollard ZF. Persistent Hyperplastic Primary Vitreous: Diagnosis, Treatment and Results. *Tr Am Ophth Soc* 1997:487-549.
- Wilson JMG, Jungner G. Principles and practice of screening for disease. *WHO Public Health Papers* 1968;34.
- Andermann A, Blancquaert I, Beauchamp S, Déry V. Revisiting Wilson and Jungner in the genomic age: a review of screening criteria over the past 40 years. *Bulletin of the World Health Organization* 2008;86(4):317-9.