

# The impact of altitude on hospitalization and hospital mortality from pandemic 2009 influenza A (H1N1) virus pneumonia in Mexico

Rogelio Pérez-Padilla, MD,<sup>(1)</sup> Cecilia García-Sancho, MD, PhD,<sup>(1)</sup> Rosario Fernández, MSc,<sup>(1)</sup> Francisco Franco-Marina, MD, MSc,<sup>(1)</sup> Hugo López-Gatell, MD, PhD,<sup>(2)</sup> Ietza Bojórquez, MD, PhD.<sup>(3)</sup>

Pérez-Padilla R, García-Sancho C, Fernández R, Franco-Marina F, López-Gatell H, Bojórquez L.  
The impact of altitude on hospitalization and hospital mortality from pandemic 2009 influenza A (H1N1) virus pneumonia in Mexico.  
*Salud Publica Mex* 2013;55:92-95.

## Abstract

**Objective.** To determine the effect of altitude of residence on influenza A (H1N1). **Materials and methods.** We analyzed 207 135 officially notified of influenza-like illness (ILI) cases, 23 048 hospitalizations and 573 deaths during the first months of the novel pandemic influenza A H1N1 virus, to examine if residents of high altitude had more frequently these adverse outcomes. **Results.** Adjusted rates for hospitalization and hospital mortality rates increased with altitude, probably due to hypoxemia.

Key words: altitude; influenza; mortality; Mexico

Pérez-Padilla R, García-Sancho C, Fernández R, Franco-Marina F, López-Gatell H, Bojórquez I.  
El impacto de la altitud en las tasas de hospitalización y mortalidad por el virus de la influenza A (H1N1) pandémico en México.  
*Salud Publica Mex* 2013;55:92-95.

## Resumen

**Objetivo.** Determinar el efecto de la altitud del lugar de residencia sobre la epidemia de influenza A (H1N1). **Materiales y métodos.** Se analizaron 207 135 casos notificados a la Secretaría de Salud, 23 048 hospitalizaciones y 573 muertes por enfermedad similar a la influenza (ILI, por sus siglas en inglés) durante los primeros meses de la pandemia del nuevo virus de la influenza A H1N1 durante 2009, para examinar si los residentes de zonas más altas presentaron tasas más altas de hospitalización y muerte. **Resultados.** Las tasas ajustadas de hospitalización y mortalidad hospitalaria se incrementaron con la altitud probablemente por la presencia de hipoxemia.

Palabras clave: altitud; influenza; mortalidad; México

(1) Instituto Nacional de Enfermedades Respiratorias (INER), México DF, México.

(2) Instituto Nacional de Salud Pública, Cuernavaca, Morelos, México.

(3) El Colegio de la Frontera Norte, Tijuana, Baja California, México.

Received on: May 9, 2012 • Accepted on: October 10, 2012

Corresponding author: Dra. Cecilia García-Sancho, Instituto Nacional de Enfermedades Respiratorias, Calzada de Tlalpan 4502, I4080 México, DF, México.

E-mail: cegarsan@netscape.net

In a previous analysis, individuals living between 2 000 and 2 499 m above sea level in Mexico had a three-fold increase in mortality due to pneumonia-influenza compared to people living below 500 m above sea level, after adjustment for poverty, age and gender.<sup>1</sup> In this study, our objective was to examine the association between altitude of residence and rates of hospitalization and mortality, in cases of Influenza-like illness (ILI) and severe acute respiratory illness (SARI), during the first months of the 2009 pandemic influenza A H1N1 virus, to examine if residents of high altitude had more frequently these adverse outcomes.

## Materials and methods

Here we analyze the first 207 135 influenza-like illness (ILI) or severe acute respiratory illness (SARI) cases, notified to the Mexican Ministry of Health (Secretaría de Salud, SSA) system from 597 sentinel health units including out-patient clinics as well as hospitals from all 32 Mexican states between June and October 2009. Notified cases came from 1 746 of the 2 445 Mexican municipalities. ILI was defined as fever ( $\geq 38^{\circ}\text{C}$ ) with cough and headache plus one additional respiratory or digestive symptom.<sup>2</sup> SARI adds to the ILI definition the presence of difficult breathing or acute respiratory failure.

Among these cases there were 23 048 hospitalizations (11%) and 573 deaths among those hospitalized (0.3%). During the same time period, an additional 24 858 cases were notified to the Health Ministry but were excluded from analysis because lacked municipality of residence. Notified cases increased progressively from May 2009, peaking at the end of September 2009 (major fall outbreak) and progressively descended in frequency during October. Laboratory confirmation, using the CDC protocol of real-time rTPCR for influenza A (H1N1), was available only for 12.5% of analyzed cases.

Several potential correlates of ILI severity were generated at the municipality level. Average altitude above the sea level of the municipalities (weighted by the population size of the communities) where cases resided was obtained from the official database of the National Institute of Statistics and Geography (Instituto Nacional de Estadística y Geografía, INEGI).<sup>3</sup> In addition, for each municipality, we obtained a 2005 normalized social deprivation index, known as the Marginality Index, with five levels, developed by the Mexican National Population Council (Consejo Nacional de Población, CONAPO) using census indicators on the educational characteristics of the resident population, the type of economic activities of the work force and the household building characteristics and access to water supply and

sewage.<sup>4</sup> Also, using the 10% population sample from the 2000 Mexican census we obtained, for each municipality of case residence, the proportion of households using wood or charcoal for cooking. Finally, data on annual mean temperatures at the municipality level were obtained from isotherms developed at CONABIO (Comisión Nacional para el Conocimiento y Uso de la Biodiversidad).<sup>5</sup>

## Statistical analysis

Multilevel logistic regression models were fitted to estimate the proportion of notified cases who were hospitalized, as well as the proportion of hospitalized cases that died (in-hospital mortality) as a function of several municipality and individual level predictors.<sup>6</sup> Municipality level predictors included in the modeling were mean altitude and annual temperature, degree of marginality and the proportion of households using wood or charcoal for cooking. Individual level predictors entered in the two fitted models were age and gender. We only evaluated mortality in hospitalized cases because in the primary care units there was not active follow up of cases and the very few deaths that were recorded occurred during the patients visit to the unit. All analyses were performed using a commercially available statistical package (Stata v.10.0).

## Results


Mean age of analyzed cases was 23.0 years (standard deviation [SD] 17.6), with 39% of them being younger than 15 years and 3.5% were 60 and older. Half (51.9%) of cases were women, 11.1% were hospitalized, 92% received treatment with oseltamivir and one in five had at least one co-morbidity recorded, mainly diabetes and asthma. Intubation as a proxy of severe acute respiratory illness (SARI) occurred in 1.4% of the total notified cases, in 16.3% of hospitalized patients and in 42% of the patients who died.

Most analyzed cases (70.0%) resided in municipalities with an average altitude lower than 1 765 m, with 37.7% residing in municipalities at <470 m and 15.0% at 2 230-2 850 m. Only 4.8% resided in municipalities with high or very high marginality. In addition, 89.3% of cases came from municipalities where less than 25% of households used wood or charcoal for cooking, another 6.4% of cases came from municipalities where between 25 and 49% of households used these fuels for cooking and only 1.3% of cases resided in municipalities where at least 75% of households used these fuels. Finally, almost all cases (94.5%) lived in municipalities with mean annual temperatures between 15°C and 27°C.

The odds of hospitalization (table I), adjusted for covariates both at the municipality level (marginality, prevalence of use of wood or charcoal for cooking in households and annual mean temperature) and at the individual level (gender and age group), were found to be higher as the altitude of residence of ILI cases increased. Adjusted hospitalization rates also increased significantly with altitude of residence (figure 1), but also in cases residing in more marginalized municipalities, in municipalities with a higher proportion of

**Table I**  
**TWO-LEVEL LOGISTIC REGRESSION MODEL OF FACTORS RELATED TO THE ODDS OF HOSPITALIZATION IN MEXICAN NOTIFIED ILI CASES**

	Odds ratio	95% Confidence interval
<i>Municipality level predictors</i>		
Altitude (meters)		
0-469	1	
470-1189	1.49	(1.37-1.62)
1190-1764	1.63	(1.52-1.74)
1765-2229	3.09	(2.83-3.38)
2230-2850	2.98	(2.71-3.28)
Marginality		
Very low	1	
Low	1.41	(1.26-1.59)
Medium	1.72	(1.54-1.92)
High	1.85	(1.48-2.31)
Very high	1.78	(1.26-2.51)
% households using wood or charcoal for cooking		
0-24	1	
25-49	1.29	(1.15-1.45)
50-74	1.54	(1.23-1.92)
75-100	1.84	(1.38-2.45)
Annual mean temperature (°C)	1.03	(1.02-1.04)
<i>Individual level predictors</i>		
Gender		
Male	1.05	(1.02-1.08)
Age group		
0-14	1	
15-29	0.805	(0.775-0.836)
30-44	1.12	(1.08-1.17)
45-59	1.75	(1.67-1.84)
60 and older	3.01	(2.84-3.20)



**FIGURE 1. THE IMPACT OF ALTITUDE ON HOSPITALIZATION FROM PANDEMIC 2009 INFLUENZA A (H1N1) VIRUS PNEUMONIA IN MEXICO. ALTITUDE ABOVE SEA LEVEL, PERCENTAGE OF HOSPITALIZED PATIENTS AND OXYGEN SATURATION ESTIMATED ACCORDING TO ALTITUDE**

households using wood or charcoal for cooking and in those with higher mean annual temperature (table I).

Table II depicts the odds of in-hospital death (adjusted by covariates) increasing with altitude of residence but reaching a plateau at the highest altitudes. Mortality was also associated to residence in a marginalized municipality but not with a higher use of wood or charcoal for cooking. Risk of hospitalization and death increased with age and in men compared to women.

## Discussion

In our analysis higher altitude of residence was associated with a higher risk of hospitalization in persons with ILI and SARI, during the 2009 influenza pandemic in Mexico in a monotonic way. Higher altitude of residence was also associated with a higher risk of death among hospitalized cases but this association seemed to reach a plateau at higher altitudes. Nevertheless, the number of deaths observed in hospitalized cases was relatively low and it is possible that we lack adequate power to evaluate the association between in-hospital mortality at the highest altitudes. Association persisted even after taking into account gender, age, and characteristics of the community such as mean ambient temperature, degree of marginality and use of biomass fuels.<sup>7</sup>

**Table II**  
**TWO-LEVEL LOGISTIC REGRESSION MODEL OF FACTORS**  
**RELATED TO THE ODDS OF DYING DURING HOSPITALIZATION**  
**IN MEXICAN NOTIFIED ILI CASES**

Municipality level predictors	Rate ratio	95% Confidence interval
Altitude (meters)		
0-469	1	
470-1189	1.80	(1.16-2.80)
1190-1764	1.61	(1.04-2.48)
1765-2229	2.37	(1.45-3.90)
2230-2850	1.45	(0.824-2.53)
Marginality		
Very low	1	
Low	0.99	(0.718-1.37)
Medium	1.06	(0.662-1.70)
High	1.31	(0.660-2.58)
Very high	1.86	(0.667-5.21)
% households using wood or charcoal for cooking		
0-24	1	
25-49	0.879	(0.531-1.45)
50-74	0.576	(0.277-1.20)
75-100	1.20	(0.503-2.86)
Annual mean temperature (°C)	0.958	(0.918-1.00)
Individual level predictors		
Gender		
Male	1.40	(1.19-1.65)
Age group		
0-14	1	
15-29	2.30	(1.68-3.14)
30-44	4.41	(3.33-5.84)
45-59	5.23	(3.93-6.96)
60 and older	5.25	(3.82-7.22)

All reported cases were part of the influenza pandemic, although the virus was detected only in 40.9% of tested cases. Negative tests are known due to inappropriate sampling and delayed testing with reduced viral loads.<sup>8</sup>

Our findings are in agreement with other reports linking altitude of residence with severe lower respiratory infections,<sup>1,9,10</sup> including influenza A H5N1<sup>11</sup> and are secondary basically to hypoxemia whose main

determinant is altitude.<sup>12</sup> A systematic review of the published literature showed that hypoxemia is a common and potentially lethal complication of acute lower respiratory infection in children less than five years of age, particularly among those with severe disease and those living at high altitude.<sup>13</sup>

According to our data in the presence of influenza and pneumonia, altitude increases the risk of severe respiratory outcomes and therefore, it is important to identify hypoxemia quickly, using an ear oximeter, and treat it with oxygen.

*Declaration of conflict of interests.* The authors declare that they have no conflict of interests.

## References

- Perez-Padilla R, Franco-Marina F. The impact of altitude on mortality from tuberculosis and pneumonia. *Int J Tuberc* 2004;8:1315-1320.
- Gupta V, Dawood FS, Rai SK, Broor S, Wigh R, Mishra AC, et al. Validity of clinical case definitions for influenza surveillance among hospitalized patients: results from a rural community in North India. *Influenza Other Respi Viruses* 2012; doi:10.1111/ij.1750-2659.2012.00401.x.
- Census of Population and Household, 2005. Main results by location (ITER) [Access date 03/20/2010]. Available at: [http://www.inegi.org.mx/sistemas/consulta\\_resultados/default.aspx](http://www.inegi.org.mx/sistemas/consulta_resultados/default.aspx)
- National Population Council. Deprivation index 2005. [Access date 03/20/2010]. Available at: <http://www.conapo.gob.mx/index.php>
- García, E. CONABIO, (1998). 'Isotermas Medias Anuales'. Escala 1:1000000, México. [Access date 03/20/2010]. Available at [http://www.conabio.gob.mx/informacion/metadatos/gis/isotm1mgw.xml?\\_httpcache=yes&\\_xsl=/db/metadatos/xsl/fgdc\\_html.xsl&\\_indent=no](http://www.conabio.gob.mx/informacion/metadatos/gis/isotm1mgw.xml?_httpcache=yes&_xsl=/db/metadatos/xsl/fgdc_html.xsl&_indent=no)
- Rahaman Khan H, Shaw EH. Multilevel Logistic Regression Analysis Applied to Binary Contraceptive Prevalence Data. *J Data Sci* 2011;9:93-110.
- Smith KR, McCracken JP, Weber MW, Hubbard A, Jenny A, Thompson LM, et al. Effect of reduction in household air pollution on childhood pneumonia in Guatemala (RESPIRE): a randomised controlled trial. *Lancet* 2011;378:1717-1726.
- CDC protocol of realtime RTPCR for influenza A (H1N1). Geneva: World Health Organization, April 2009. [Access date 03/20/2010] Available at: <http://www.who.int/csr/resources/publications/swineflu/real-timeptpcr/en/index.html>
- Khan AJ, Hussain H, Omer SB, Chaudry S, Ali S, Khan A, et al. High incidence of childhood pneumonia at high altitudes in Pakistan: a longitudinal cohort study. *Bull World Health Organ* 2009;87:193-199.
- Choudhuri JA, Ogden LG, Ruttenber AJ, Thomas DS, Todd JK, Simoes EA. Effect of altitude on hospitalizations for respiratory syncytial virus infection. *Pediatrics* 2006; 117:349-356.
- Ge E, Haining R, Li CP, Yu Z, Wayne MY, Chu KH et al. Using knowledge fusion to analyze avian influenza H5N1 in East and Southeast Asia. *PLoS One* 2012;7(5):e29617.
- Perez-Padilla R, Torre-Bouscoulet L, Muino A, Marquez MN, Lopez MV, de Oca MM, et al. Prevalence of oxygen desaturation and use of oxygen at home in adults at sea level and at moderate altitude. *Eur Respir J* 2006;27:594-599.
- Lozano JM. Epidemiology of hypoxaemia in children with acute lower respiratory infection. *Int J Tuberc Lung Dis* 2001;5:496-504.